

ON THE ARROW PROPERTY  
SH782

SAHARON SHELAH

arXiv:math/0112213v1 [math.LO] 20 Dec 2001

---

I would like to thank Alice Leonhardt for the beautiful typing.  
Done 2/4/01  
Latest Revision - 01/Dec/15

Typeset by  $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{\textsf{TeX}}$

## §0 INTRODUCTION

Let  $X$  be a finite set of alternatives. A choice function  $c$  is a mapping which assigns to nonempty subsets  $S$  of  $X$  an element  $c(S)$  of  $S$ . A *rational* choice function is one for which there is a linear ordering on the alternatives such that  $c(S)$  is the maximal element of  $S$  according to that ordering. (We will consider choice functions which are defined on subsets of  $X$  of fixed cardinality  $k$ .)

Arrow's impossibility theorem [Arr50] asserts that under certain natural conditions, if there are at least three alternatives then every non-dictatorial social choice gives rise to a non-rational choice function, i.e., there exist profiles such that the social choice is not rational. A profile is a finite list of linear orders on the alternatives which represent the individual choices. For general references on Arrow's theorem and social choice functions see [Fis73], [Pel84] and [Sen86].

Non-rational classes of choice functions which may represent individual behavior where considered in [KRS01] and [Kal01]. For example:  $c(S)$  is the second largest element in  $S$  according to some ordering, or  $c(S)$  is the median element of  $S$  (assume  $|S|$  is odd) according to some ordering. Note that the classes of choice functions in these classes are *symmetric* namely are invariant under permutations of the alternatives. Gil Kalai asked if Arrow's theorem can be extended to the case when the individual choices are not rational but rather belong to an arbitrary non-trivial symmetric class of choice functions. (A class is non-trivial if it does not contain all choice functions.) The main theorem of this paper gives an affirmative answer in a very general setting. See also [RF86] for general forms of Arrow's and related theorem.

The main result is related to clones which are studied in universal algebras (but we do not use this theory). On clones see [Szb99] and [Szn96].

Notation:

- 1)  $n, m, k, \ell, r, s, t, i, j$  natural numbers always  $k$ , many times  $r$  are constant (there may be some misuses of  $k$ ).
- 2)  $X$  a finite set.
- 3)  $\mathfrak{C}$  a family of choice function on  $\binom{X}{k} = \{Y : Y \subseteq X, |Y| = k\}$ .
- 4)  $\mathcal{F}$  is a clone on  $X$  (see Definition 2.3(2)).
- 5)  $a, b, e \in X$ .
- 6)  $c, d, \in \mathfrak{C}$ .
- 7)  $f, g \in \mathcal{F}$ .

## ANNOTATED CONTENT

## §1 Framework

[What are  $X, \mathfrak{C}, \mathcal{F} = Av(\mathfrak{C})$ , the Arrow property restriction to  $\binom{X}{k}$ ,  $\mathfrak{C}$  is  $(X, k) = FCF$  (debt: no connection for different  $k - S$ ) and the Main theorem. For  $\mathfrak{C}, \mathcal{F}, r = r(\mathcal{F})$ .]

Part A: The simple case.

## §2

[Define a clone,  $r(\mathcal{F})$ . If  $f \in \mathcal{F}_{(r)}$  is not a monarchy,  $r \geq 4$  on the family of not one-to-one sequences  $\bar{a} \in {}^r X$  then  $f$  is a projection, 2.5.

Define  $f_{r;\ell,k}$ , basic implications on  $f_{r;\ell,k} \in \mathcal{F}$ , 2.6, 2.7.

If  $r = 3, f \in \mathcal{F}_{[s]}$  is not a monarchy on one-to-one triples, then  $f$  without loss of generality it is  $f_{r;1,2}$  or  $g_{r;1,2}$  on then 2.8.

If  $r = 3, f$  is not a semi monarchy on permutations of  $\bar{a}$ .

If  $r = 3$ , there are some “useful”  $f$ , 3.1. Implications on  $f_{r;\ell,k} \in \mathcal{F}$ .]

§3 Getting  $\mathfrak{C}$  is full

[Sufficient condition for  $r \geq 4$  with  $f_{r;1,2}$  or so (3.1), end?  $r = 3$ ?

Sufficient condition for  $r = 3$  with  $g_{r;1,2}$  or so (3.3).

A pure sufficient condition for  $\mathfrak{C}$  full 3.4.

Subset  $\binom{X}{3}$ , closed under a distance 3.5.

Getting the final conclusion (relying on 3, 3.6).]

§4 The  $r = 2$  case.

[By stages we get a  $f \in \mathcal{F}_{[r]}$  which is a monarchy with exactly one exceptional pair 4.2 - 4.4. Then by composition we get  $g \in \mathcal{F}_2$  similar to  $f_{r;1,2}$ .]

Part B: Non simple case.

[Maybe we can start with this and prove from part A only what is used; important question from presentation does not interest me.]

## §5 Fullness - the nonsimple case

[We derive “ $\mathfrak{C}$  is full” from various assumptions, and then prove the main theorem.]

§6 The case  $r = 2$ .§7 The case  $r \geq 4$ .

## §1 FRAMEWORK

**1.1 Context.** We fix a finite  $X$  and  $r = \{0, \dots, r-1\}$ .

**1.2 Definition.** 1) An  $(X, r)$ -election rule is a function  $c$  such that: for every “vote”  $\bar{t} = \langle t_a : a \in X \rangle \in {}^X r$  we have  $c(\bar{t}) \in r = \{0, \dots, r-1\}$ .

2)  $c$  is a monarchy if  $(\exists a \in X)(\forall \bar{t} \in {}^X \bar{r})[c(\bar{t}) = t_a]$ .

3)  $c$  is reasonable if  $(\forall \bar{t})(c(\bar{t}) \in \{t_a : a \in X\})$ .

**1.3 Definition.** 1) We say  $\mathfrak{C}$  is a family of choice functions for  $X$  ( $X$ -FCF in short) if:

$\mathfrak{C} \subseteq \{c : c \text{ is a function}$

$\text{Dom}(c) = \mathcal{P}^-(X) (= \text{family of nonempty subsets of } X)$   
and  $(\forall Y \in \mathcal{P}^-(X))(c(Y) \in Y)\}$ .

2)  $\mathfrak{C}$  is called symmetric if for every  $\pi \in \text{Per}(X) = \text{group of permutations of } X$  we have

$$c \in \mathfrak{C} \Rightarrow \pi * c \in \mathfrak{C}$$

where

$$\pi * c(Y) = \pi^{-1}(c(\pi(Y))).$$

3)  $\mathcal{P}_{\mathfrak{C}} = \mathcal{P}^-(X)$ .

**1.4 Definition.** 1) We say  $\text{av}$  is a  $r$ -averaging function for  $\mathfrak{C}$  if

(a)  $\text{av}$  is a function written  $\text{av}_Y(a_1, \dots, a_r)$

(b) for any  $c_1, \dots, c_r \in \mathfrak{C}$ , there is  $c \in \mathfrak{C}$  such that  
 $(\forall Y \in \mathcal{P}^-(X))(c(Y)) = \text{av}_Y(c_1(Y), \dots, c_r(Y))$

(c) if  $a \in Y \in \mathcal{P}^-(X)$  then  $\text{av}_Y(a, \dots, a) = a$ .

2)  $\text{av}$  is simple if  $\text{av}_Y(a_1, \dots, a_r)$  does not depend on  $Y$  so we may omit  $Y$ .

3)  $\text{AV}_r(\mathfrak{C}) = \{\text{av} : \text{av is an } r\text{-averaging function for } \mathfrak{C}\}$ , similarly  $\text{AV}_r^s(\mathfrak{C}) = \{\text{av} : \text{av is a simple } r\text{-averaging function for } \mathfrak{C}\}$ .

4)  $\text{AV}(\mathfrak{C}) = \bigcup_r \text{AV}_r(\mathfrak{C})$  and  $\text{AV}^s(\mathfrak{C}) = \bigcup_r \text{AV}_r^s(\mathfrak{C})$ .

**1.5 Definition.** 1)  $\mathfrak{C}$  has the simple  $r$ -Arrow property if

$$\text{av} \in AV_r^s(\mathfrak{C}) \Rightarrow \bigvee_{t=1}^r (\forall a_1, \dots, a_2) \text{av}(a_1, \dots, a_2) = a_t$$

such  $\text{av}$  is called monarchical.

2) Similarly without simple.

1.6 Question: 1) When does  $\mathfrak{C}$  have the Arrow property?

2) Does  $|\mathfrak{C}| \leq \text{poly}(|X|) \Rightarrow r$ -Arrow property?

*1.7 Remark.* The question was asked with  $\mathfrak{C}_{(X)}$  defined for every  $X$ ; but in the treatment this does not influence.

We actually deal with:

**1.8 Definition.** If  $1 \leq k \leq |X| - 1$  and we replace  $\mathcal{P}^-(X)$  by  $\binom{X}{k} =: \{Y : Y \subseteq X, |Y| = k\}$ , then  $\mathfrak{C}$  is called  $(X, k)$  - FCF,  $\mathcal{P}_{\mathfrak{C}} = \binom{X}{k}, k = k(\mathfrak{C})$ ,  $\text{av}$  is [simple]  $r$ -averaging function for  $\mathfrak{C}$ ; let  $k(\mathfrak{C}) = \infty$  if  $\mathcal{P}_{\mathfrak{C}} = \mathcal{P}^-(X)$ ; let  $\mathcal{F}(\mathfrak{C}) = Av^s(\mathfrak{C})$  let  $\mathcal{F}_{[r]} = \{f \in \mathcal{F} : f \text{ is } r\text{-place}\}$ .

1.9 Discussion: This is justified:

1) For simple averaging function,  $k \geq r$  the restriction to  $\binom{X}{k}$  implies the full result. [In the end, draw the conclusion.]

2) For the nonsimple case there is a little connection between the various  $\mathfrak{C} \upharpoonright \binom{X}{k}$  (exercise). Our aim is (but we shall first prove the simple case):

**1.10 Main Theorem.** *There are  $r_1^*, r_2^* < \omega$ . (We shall be able to give explicit values, e.g.  $r_1^* = r_2^* = 7$  are O.K.) such that:*

- ⊗ *if  $X$  is finite,  $r_1^* \leq k, |X| - r_2^* \geq k$  and  $\mathfrak{C}$  is an  $(X, k)$ -FDF and some  $\text{av} \in AV_r(\mathfrak{C})$  is not monarchical, then every choice function for  $\binom{X}{k}$  belongs to  $\mathfrak{C}$  ( $= \mathfrak{C}$  is full).*

Part A: Simple case.

## §2 CONTEXT AND ON NICE $f$ 'S

Note: Sometimes Part B gives alternative ways.

*2.1 Hypothesis (for part A).*

- (a)  $X$  a finite set
- (b)  $5 < k < |X| - 5$
- (c)  $\mathfrak{C}$  a symmetric  $(X, k)$ -FCF and  $\mathfrak{C} \neq \emptyset$
- (d)  $\mathcal{F}_{[r]} = \{f : f \text{ an } r\text{-place function from } X \text{ to } X \text{ such that } \mathfrak{C} \text{ is closed under } f \text{ that is } f \in \text{AV}_r^s(\mathfrak{C})\}$
- (e)  $\mathcal{F} = \cup\{\mathcal{F}_{[r]} : r < \omega\}$
- (f)  $r(\mathcal{F}) = \text{Min}\{r : \text{some } f \in \mathcal{F}_{[r]} \text{ is not a monarchy}\}.$

*2.2 Fact.*  $\mathcal{F}$  is a clone on  $X$  (see 2.3 below).

**2.3 Definition.** 1)  $f$  is monarchical = projection if  $f$  is an  $r$ -place function (from  $X$  to  $X$ ) and for some  $t$ ,  $(\forall x_1, \dots, x_n) f(x_1, \dots, x_r) = x_t$ .  
 2)  $\mathcal{F}$  is a clone on  $X$  if it is a family of functions from  $X$  to  $X$  (for all arities) including the projections and closed under composition.

**2.4 Definition.** For  $\mathfrak{C}, \mathcal{F}$  as in 2.1:

$$r(\mathfrak{C}) = r_m(\mathcal{F}) =: \text{Min}\{r : \text{some } f \in \mathfrak{C}_r \text{ is not monarchical}\}$$

(let  $r(\mathcal{F}) = \infty$  if  $\mathfrak{C}$  is monarchical).

**2.5 Claim.** Assume

- (a)  $f \in \mathcal{F}_{[r]}$
- (b)  $4 \leq r = r_m(\mathcal{F}) = \text{Min}\{r : \text{some } f \in \mathcal{F} \text{ is not a monarchy}\}.$

Then

- 1) for some  $\ell \in \{1, \dots, r\}$  we have  $f(x_1, \dots, x_r) = x_\ell$  if  $x_1, \dots, x_r$  has some repetition.  
 2)  $r \leq k$ .

*Proof.* 1) Clearly there is a two-place function  $h$  from  $\{1, \dots, r\}$  to  $\{1, \dots, r\}$  such that: if  $y_\ell = y_k \wedge \ell \neq k \Rightarrow f(y_1, \dots, y_r) = y_{h(\ell, k)}$ ; we have some freedom so let without loss of generality

$$\boxtimes \ell \neq k \Rightarrow h(\ell, k) \neq k.$$

Assume toward contradiction that

$$\circledast (1)'s \text{ conclusion fails, i.e. } h \upharpoonright h(\ell, k) : 1 \leq \ell < k \leq r \text{ is not constant.}$$

Case 1: For some  $\bar{x} \in {}^r X$  and  $\ell_1 \neq k_1 \in \{1, \dots, r\}$  we have

$$x_{\ell_1} = x_{k_1}$$

$$f(\bar{x}) \neq x_{\ell_1}$$

equivalently:  $h\{\ell_1, k_1\} \notin \{\ell_1, k_1\}$ .

Without loss of generality  $\ell_1 = r - 1, k_1 = r, f(\bar{x}) = x_1$  (as for a permutation  $\sigma$  of  $\{1, \dots, r\}$  we can replace  $f$  by  $f_\sigma, f_\sigma(x_1, \dots, x_r) = f(x_{\sigma(1)}, \dots, x_{\sigma(r)})$ ).

We can choose  $x \neq y$  in  $X$  so  $h(x, y, \dots, y) = x$  hence  $\ell \neq k \in \{2, \dots, r\} \Rightarrow h(\ell, k) = 1$ .

Now for  $\ell \in \{2, \dots, r\}$  we have agreed  $h(1, \ell) \neq \ell$ , (see  $\boxtimes$ ) so as  $h \upharpoonright \{(\ell, k) : \ell < k\}$  is not constantly 1 (by  $\circledast$ ) without loss of generality  $h(1, 2) = 3$ . But as  $r \geq 4$  letting  $x \neq y \in X$  we have  $f(x, x, y, y)$  is  $y$  as  $h(1, 2) = 3$  and is  $x$  as  $h(3, 4) = 1$ , contradiction.

Case 2: Not Case 1.

Let  $x \neq y$  compute  $f(x, x, y, y, \dots)$  it is  $x$  as  $h(1, 2) \in \{1, 2\}$  and it is  $y$  as  $h(3, 4) \in \{3, 4\}$ , contradiction.

2) Follows as for  $r > k$  we always have a repetition.  $\square_{2.5}$

**2.6 Definition.**  $f_{r;\ell,k} = f_{r,\ell,k}$  is the  $r$ -place function on  $X$  defined by

$$f_{r;\ell,k}(\bar{x}) = \begin{cases} x_\ell & \bar{x} \text{ is with repetition} \\ x_k & \text{otherwise} \end{cases}$$

**2.7 Claim.** 1) If  $f_{r,1,2} \in \mathcal{F}$  then  $f_{r,\ell,k} \in \mathfrak{C}$  for  $\ell \neq k \in \{1, \dots, r\}$ .  
 2) If  $f_{r,1,2} \in \mathcal{F}$  and  $r = r[\mathcal{F}] \geq 3$  then  $f_{r+1,1,2} \in \mathcal{F}$ .

*Proof.* 1) Trivial.

2) First assume  $r > 5$ . Let  $g(x_1, \dots, x_{r+1}) = f_{r,1,2}(x_1, x_2, \tau_3 \dots \tau_r)$  where  $\tau_m \equiv f_{r,1,m}(x_1, \dots, x_m, x_{m+2}, \dots, x_{r+1})$ ; (that is  $x_{m+1}$  is omitted).

So for any  $\bar{a}$ :

if  $\bar{a}$  with no repetitions:

$$\tau_3(\bar{a}) = a_3, \dots, \tau_m(\bar{a}) = a_m$$

$$g(\bar{a}) = f(a_1, a_2, a_3, \dots, a_r) = a_2$$

if  $\bar{a}$  has repetitions say  $a_\ell = a_k$  then there is  $m \in \{3, \dots, r\} \setminus \{\ell - 1, k - 1\}$  hence  $\langle a_1, \dots, a_m, a_{m+2}, \dots, a_{r+1} \rangle$  is with repetition so  $\tau_m(\bar{a}) = a_1$  so  $(a_1, a_2, \dots, \tau_m(\bar{a}), \dots)$  has repetition so  $g(\bar{a}) = a_1$ .

Second assume  $r = 4$ :

Let  $(\tau_1, \dots, \tau_4), g$  be of arity 5 and for  $\bar{x} = (x_1, \dots, x_5)$  we let  $g(\bar{x}) = f_{r,1,2}(\tau_1(\bar{x}), \dots, \tau_4(\bar{x}))$  where

$$(*)_1 \quad \tau_1(\bar{x}) = x_1$$

$$(*)_2 \quad \tau_2(\bar{x}) = f_{r,1,2}(x_1, x_2, x_3, x_4)$$

$$(*)_3 \quad \tau_3(\bar{x}) = f_{r,1,3}(x_1, x_2, x_3, x_5), \tau_4(\bar{x}) = f_{r,1,3}(x_1, x_2, x_4, x_5).$$

Note that

$$(*)_4 \quad \text{for } \bar{x} \text{ with no repetition } \tau_\ell(\bar{x}) = x_\ell.$$

Now check that  $g$ , i.e. is as required.

Third assume  $r = 3$ :

Let  $g'(x_1, x_2, x_3, x_4) = f_{r,1,2}(\tau_1, \tau_2, \tau_3)$  where

$$\tau_1 = x_1$$

$$\tau_2 = f_{r,1,2}(x_1, x_2, x_4)$$

$$\tau_3 = f_{r,1,2}(x_2, x_3, x_4).$$

Now check (or see 4.7's proof).

□<sub>2.7</sub>



**2.8 Claim.** *Assume*

- ( $\alpha$ )  $\mathcal{F}$  is a symmetric clone,
- ( $\beta$ ) every  $f \in \mathcal{F}_{[2]}$  is a monarchy,  $r = r[\mathcal{F}] = 3$
- ( $\gamma$ )  $f^* \in \mathcal{F}_{[3]}$  and for no  $i \in \{1, 2, 3\}$  do we have  $(\forall \bar{b} \in {}^3X)(\bar{b} \text{ not one-to-one} \Rightarrow f^*(\bar{b}) = b_i)$ .

Then for some  $g \in \mathcal{F}_{[3]}$  not a monarchy we have: (a) and (b) where

- (a) for  $\bar{b} \in {}^3X$  which is not one-to-one  $g(\bar{b}) = f_{r;1,2}(\bar{b})$ , i.e.  $= b_1$
- (b) for  $\bar{b} \in {}^3X$  which is not one-to-one  $g(\bar{b}) = g_{r;1,2}(\bar{b})$ , see below

Where

**2.9 Definition.**  $g_{r;1,2}$  is the following function<sup>1</sup> from  $X$  to  $X$ .

$$g_{r;1,2}(x_1, x_2, \dots, x_r) = \begin{cases} x_2 & \text{if } x_2 = x_3 = \dots = x_r \\ x_1 & \text{if otherwise} \end{cases}.$$

*Proof of 2.8.* The same as the proof of the next claim ignoring the one-to-one sequences (i.e.  $f(a_1, a_2, a_3)$ ), see more later.

**2.10 Claim.** *Assume  $\mathcal{F}$  is a symmetric clone,  $r = r(\mathcal{F}) = 3$ ,  $f^* \in \mathcal{F}$ ,  $f^*$  is a 3-place function and not a monarchy and  $\bar{a} \in {}^3X$  is with no repetition such that: if  $\bar{a}' = (a'_1, a'_2, a'_3)$  is a permutation of  $\bar{a}$  then  $f^*(\bar{a}') = a'_1$ ; but  $\neg(\forall \bar{b} \in {}^3X)(\bar{b} \text{ not one-to-one} \rightarrow f(\bar{b}) = b_1)$ .*

Then for some  $g \in \mathcal{F}_3$  we have (a) or we have (b) where:

- (a)(i) for  $\bar{b} \in {}^3X$  with repetition,  $g(\bar{b}) = f_{r;1,2}(\bar{b})$ , i.e.  $g(\bar{b}) = b_1$
- (ii)  $g(\bar{a}') = a'_2$  for any permutation  $\bar{a}'$  of  $\bar{a}$ ,  $\bar{b}$
- (b)(i) for  $\bar{b} \in {}^3X$  with repetition,  $g(\bar{b}) = g_{r;1,2}(\bar{b})$
- (ii)  $g(\bar{a}') = a'_1$  for any permutation  $\bar{a}'$  of  $\bar{a}$  (see on  $g_{r;1,2}$  in 2.9).

*Proof.* Let  $\bar{a} = (a_1, a_2, a_3)$ ;  $(a, b, c)$  denote any permutation of  $\bar{a}$ .

Let  $W = \{\bar{b} : \bar{b} \in {}^3X \text{ and } [\bar{b} \text{ is a permutation of } \bar{a} \text{ or } \bar{b} \text{ not one-to-one}]\}$ .  
 Let  $\mathcal{F}^- = \{f \upharpoonright W : f \in \mathcal{F}\}$ ,  $f = f^* \upharpoonright W$ ; recall that  $f(a_1, a_2, a_3) = a_1$ .  
 Let for  $\eta \in \{1, 2\}$ ,  $f_\eta$  be the 3-place function with domain  $W$ , such that [remember that  $f(x, y, y), f(x, y, x), f(x, x, y)$  are monarchies and]:

---

<sup>1</sup>this is the majority function for  $r = 3$

- $\boxtimes_0 f_\eta(a_{\sigma(1)}, a_{\sigma(2)}, a_{\sigma(3)}) = a_{\sigma(1)}$  for  $\sigma \in \text{Per}\{1, 2, 3\}$
- $\boxtimes_1 f_\eta(a_1, a_2, a_2) = a_{\eta(1)}$
- $\boxtimes_2 f_\eta(a_1, a_2, a_1) = a_{\eta(2)}$
- $\boxtimes_3 f_\eta(a_1, a_1, a_2) = a_{\eta(3)}$

Now

- $(*)_0 f \in \{f_\eta : \eta \in {}^3 2\}$   
[why? just think: by the assumption on  $f^*$  and as  $r(\mathcal{F}) = 3$ ]
- $(*)_1$  if  $\eta = \langle 1, 1, 1 \rangle$  then  $f_\eta$  is  $\neq f$   
[why?  $f_\eta(x_1, x_2, x_3) = x_1$  on  $W$ , i.e. is a monarchy]
- $(*)_2$  if  $\eta, \nu \in {}^3\{1, 2\}$ ,  $\eta(1) = \nu(1), \eta(2) = \nu(3), \eta(3) = \nu(2)$ , then  $f_\eta \in \mathcal{F}^- \Leftrightarrow f_\nu \in \mathcal{F}^-$   
[Why? In  $f(x, y, z)$  we just exchange  $y$  and  $z$ ]
- $(*)_3$  if  $f_{\langle 2, 2, 2 \rangle} \in \mathcal{F}^-$  then  $f_{\langle 1, 2, 2 \rangle} \in \mathcal{F}^-$   
[Why? Define  $g$  by  $g(x, y, z) = f_{\langle 2, 2, 2 \rangle}(x, f_{\langle 2, 2, 2 \rangle}(y, x, z), f_{\langle 2, 2, 2 \rangle}(z, x, y))$   
(so  $g \in \mathcal{F}^-$ ) hence

$$g(a, b, c) = f_{\langle 2, 2, 2 \rangle}(a, b, c), \text{ generally } g \text{ satisfies } \boxtimes_0$$

$$\begin{aligned} g(a, b, b) &= f_{\langle 2, 2, 2 \rangle}(a, f_{\langle 2, 2, 2 \rangle}(b, a, b), f_{\langle 2, 2, 2 \rangle}(b, a, b)) \\ &= f_{\langle 2, 2, 2 \rangle}(a, a, a) = a \end{aligned}$$

$$\begin{aligned} g(a, b, a) &= f_{\langle 2, 2, 2 \rangle}(a, f_{\langle 2, 2, 2 \rangle}(b, a, a), f_{\langle 2, 2, 2 \rangle}(a, a, b)) \\ &= f_{\langle 2, 2, 2 \rangle}(a, a, b) = b \end{aligned}$$

$$\begin{aligned} g(a, a, b) &= f_{\langle 2, 2, 2 \rangle}(a, f_{\langle 2, 2, 2 \rangle}(a, a, b), f_{\langle 2, 2, 2 \rangle}(b, a, a)) \\ &= f_{\langle 2, 2, 2 \rangle}(a, b, a) = b. \end{aligned}$$

- So  $g = f_{\langle 1, 2, 2 \rangle}$
- $(*)_4 f_{\langle 1, 2, 2 \rangle} \in \mathcal{F}^- \Rightarrow f_{\langle 2, 1, 2 \rangle} \in \mathcal{F}^-$   
[Why? Let

$$g(x, y, z) = f_{<1,2,2>}(x, y, f_{<1,2,2>}(z, x, y))$$

So

$$g(a, b, c) = a \text{ and } g \text{ satisfies } \boxtimes_0$$

and

$$\begin{aligned} g(a, b, b) &= f_{<1,2,2>}(a, b, f_{<1,2,2>}(b, a, b)) \\ &= f_{<1,2,2>}(a, b, a) = b \end{aligned}$$

$$\begin{aligned} g(a, b, a) &= f_{<1,2,2>}(a, b, f_{<1,2,2>}(a, a, b)) \\ &= f_{<1,2,2>}(a, b, b) = a \end{aligned}$$

$$\begin{aligned} g(a, a, b) &= f_{<1,2,2>}(a, a, f_{<1,2,2>}(b, a, a)) \\ &= f_{<1,2,2>}(a, a, b) = b. \end{aligned}$$

So  $g = f_{<2,1,2>}$  so  $f_{<2,1,2>} \in \mathcal{F}^-$  as promised

(\*)<sub>5</sub>  $f_{<2,1,2>} = f_{3;3,1}$ , i.e.

$$f_{<2,1,2>}(x_1, x_2, x_3) = \begin{cases} x_1 & \underline{\text{if}} \quad |\{x_1, x_2, x_3\}| = 3 \text{ (i.e. } \{x_1, x_2, x_3\} = \bar{a}) \\ x_3 & \underline{\text{if}} \quad |\{x_1, x_2, x_3\}| \leq 2 \end{cases}$$

when  $(x_1, x_2, x_3) \in W$

[Why? Check.]

(\*)<sub>6</sub>  $f_{<2,2,1>}(x_1, x_2, x_3) = x_2$  if  $2 \geq |\{x_1, x_2, x_3\}|$   
[Why? Check.]

(\*)<sub>7</sub>  $f_{<2,1,2>} \in \mathcal{F}^- \Leftrightarrow f_{<2,2,1>} \in \mathcal{F}^-$   
[Why? See (\*)<sub>2</sub> in the beginning.]

(\*)<sub>8</sub>  $f_{<1,2,1>} \in \mathcal{F}^- \Leftrightarrow f_{<1,1,2>} \in \mathcal{F}^-$   
[Why? By (\*)<sub>2</sub> in the beginning.]

(\*)<sub>9</sub>  $f_{<1,2,1>} \in \mathcal{F}^- \Rightarrow f_{<2,2,1>} \in \mathcal{F}^-.$

Let  $g(x, y, z) = f_{<1,2,1>}(x, f_{<1,2,1>}(y, z, x), f_{<1,2,1>}(z, x, y))$

$$\begin{aligned}
g(a, b, c) &= f_{\langle 1, 2, 1 \rangle}(a, f_{\langle 1, 2, 1 \rangle}(b, c, a), f_{\langle 1, 2, 1 \rangle}(c, a, b)) \\
&= f_{\langle 1, 2, 1 \rangle}(a, b, c) = a \\
&\text{and generally } g \text{ satisfies } \boxtimes_0
\end{aligned}$$

$$g(a, b, b) = f_{\langle 1, 2, 1 \rangle}(a, f_{\langle 1, 2, 1 \rangle}(b, b, a), f_{\langle 1, 2, 1 \rangle}(b, a, b)) = f_{\langle 1, 2, 1 \rangle}(a, b, a) = b$$

$$g(a, b, a) = f_{\langle 1, 2, 1 \rangle}(a, f_{\langle 1, 2, 1 \rangle}(b, a, a), f_{\langle 1, 2, 1 \rangle}(a, a, b)) = f_{\langle 1, 2, 1 \rangle}(a, b, a) = b$$

$$g(a, a, b) = f_{\langle 1, 2, 1 \rangle}(a, f_{\langle 1, 2, 1 \rangle}(a, b, a), f_{\langle 1, 2, 1 \rangle}(b, a, a)) = f_{\langle 1, 2, 1 \rangle}(a, b, b) = a.$$

So  $g = f_{\langle 2, 2, 1 \rangle}$  hence  $f_{\langle 2, 2, 1 \rangle} \in \mathcal{F}^-$ .

## DIAGRAM

Diagram (arrows mean belonging to  $\mathcal{F}^-$  follows)

$$\begin{array}{ccc}
 & f_{\langle 2,2,2 \rangle} \in \mathcal{F}^- & \\
 & \downarrow (*)_3 & \\
 f_{\langle 1,2,2 \rangle} \in \mathcal{F}^- & & f_{\langle 1,2,1 \rangle} \in \mathcal{F}_{(*)_8}^- \xleftrightarrow{(*)_8} \mathcal{F}_{\langle 1,1,2 \rangle} \in \mathcal{F}^- \\
 & \downarrow (*)_4 & (*)_9 \swarrow \\
 & f_{\langle 2,1,2 \rangle} \in \mathcal{F} \xleftrightarrow{(*)_7} f_{\langle 2,2,1 \rangle} \in \mathcal{F}^- & 
 \end{array}$$

among the  $2^3$  function  $f_\eta$  one is discarded being a monarchy, see  $(*)_1$ , six appear in the diagram above and implies  $f_{r;2,1} \in \mathcal{F}^-$  by  $(*)_5$  hence clause (a) holds, and one is  $g_{r;1,2}$  because

$$(*) \ g_{r;1,2} = f_{\langle 2,1,1 \rangle} \text{ on } W.$$

(Why check?), so clause (b) holds. □<sub>2.10</sub>

Continuation of the proof of ?: As  $r(\mathcal{F}) = 3$  for some  $\eta \in {}^3 2$ ,  $f^*$  agrees with  $f_\eta$  for all not one-to-one triples  $\bar{b}$ . If  $\eta = \langle 1, 1, 1 \rangle$  we contradict an assumption and if  $\eta = \langle 2, 1, 1 \rangle$ , possibility (b) of 2.8 holds. If  $\eta = \langle 2, 1, 2 \rangle$  then  $f^*(\bar{b}) = b_2$  for  $\bar{b} \in {}^3 X$  not one-to-one (see  $(*)_5$ ) and as  $f^*$  is not a monarchy we get possibility (a), possibly permutating of the variables; similarly  $\eta = \langle 2, 2, 1 \rangle$ . In the remaining case (see the diagram), there is  $f \in \mathcal{F}$  agreeing on  $\{\bar{b} \in {}^3 X : \bar{b} \text{ is not one-to-one}\}$  with  $f_\eta$  for  $\eta = \langle 1, 2, 2 \rangle$  or  $\eta = \langle 1, 2, 1 \rangle$ , without loss of generality  $f^* = f$ .

If  $\eta = \langle 1, 2, 2 \rangle$  define  $g$  as in  $(*)_4$ , i.e.  $g(x, y, z) = f^*(x, y, f^*(z, x, y))$  so for a non one-to-one sequence  $\bar{b} \in {}^3 X$  we have  $g(\bar{b}) = f_{\langle 2,1,2 \rangle}(\bar{b}) = b_3$ . If for some one-to-one  $\bar{a} \in {}^3 X$  we have  $f^*(a_3, a_1, a_2) \neq a_3$  then  $g(a_1, a_2, a_3) = f^*(a_1, a_2, f^*(a_3, a_1, a_2)) \in \{a_1, a_2\}$  so permuting the variables we get possibility (a). So we are left with the case  $\bar{a} \in {}^3 X$  is one-to-one  $\Rightarrow f^*(\bar{a}) = a_1$ .

Let us define  $g \in \mathcal{F}_{[3]}$  by  $g(x_1, x_2, x_3) = f^*(f(x_2, x_3, x_1), x_3, x_2)$ . Let  $\bar{b} \in {}^3 X$ ; if  $\bar{b}$  is with no repetitions then  $g(\bar{b}) = f^*(b_2, b_3, b_2) = b_3$ . If  $\bar{b} = (a, b, b)$  then  $g(\bar{b}) = f^*(f^*(b, b, a), b, b) = f^*(a, b, b) = a = b_1$  and if  $\bar{b} = (a, b, a)$  then  $g(\bar{b}) = f^*(f^*(b, a, a), a, b) = f^*(b, a, b) = a = b_1$  and if  $\bar{b} = (a, a, b)$  then  $g(\bar{b}) = f^*(f^*(a, a, b), b, a) = f^*(b, b, a) = a = b_1$ , so for  $\bar{b}$  not one to one,  $g(\bar{b}) = b_1$ . So  $g$  is as required in clause (a).

Lastly, let  $\eta = \langle 1, 2, 1 \rangle$  and let  $g(x, y, z) = f^*(x, f^*(y, z, x), f^*(z, x, y))$  by  $(*)_9$ , easily  $[\bar{b}$  is not one-to-one  $\Rightarrow g(\bar{b}) = f_{\langle 2, 2, 1 \rangle}(\bar{b}) = b_2]$ . Now  $(a_1, a_2, a_3)$  is with no repetitions and  $f^*(a_2, a_3, a_1) = a_1$  then  $g(a_1, a_2, a_3) = a_1$  and possibility (a) holds. Otherwise for  $[\bar{b} \in {}^3X$  is one-to-one  $\Rightarrow f^*(\bar{b}) \in \{b_1, b_2\}]$ ; so if  $(a_1, a_2, a_3) \in {}^3X$  is one-to-one and  $f^*(a_2, a_3, a_1) \neq a_2$  then  $g(a_1, a_2, a_3) \neq a_2$  (as  $f^*(a_3, a_1, a_2) \neq a_2$ ) so possibility (a) holds. Hence  $[\bar{b} \in {}^3X$  is one-to-one  $\Rightarrow f^*(\bar{b}) = b_2]$ . Let  $g^*$  be  $g^*(x, y, z) = f^*(f^*(x, y, z), f^*(x, z, y), x)$ . Now if  $\bar{b}$  is one to one then  $g^*(\bar{b}) = f^*(b_2, b_3, b_1) = b_3$ . Also if  $\bar{b} = (a, b, b)$  then  $g^*(\bar{b}) = f^*(f^*(a, b, b), f^*(a, b, b), a) = f^*(a, a, a) = a$ , and if  $\bar{b} = (a, b, a)$  then  $g^*(\bar{b}) = f^*(f^*(a, b, a), f^*(a, a, b), a) = f^*(b, a, a) = b$  and if  $\bar{b} = (a, a, b)$  then  $g^*(\bar{b}) = f^*(f^*(a, a, b), f^*(a, b, a), a) = f^*(a, b, a) = b$ . So  $g^*$  is as required in the case  $\eta = \langle 1, 2, 2 \rangle$ .  $\square_{2.8}$

### 3.1 Claim. Assume

- ( $\alpha$ )  $\mathcal{F}$  is a symmetric clone of  $X$
- ( $\beta$ ) every  $f \in \mathcal{F}_{[2]}$  is monarchical
- ( $\gamma$ )  $f^* \in \mathcal{F}_{[3]}$  is not monarchical.

Then one of the following holds

- (a) for every one-to-one  $\bar{a} \in {}^3X$  for some  $f = f_{\bar{a}}$  we have
  - (i)  $f_{\bar{a}}(\bar{a}) = a_2$  (even  $f_{\bar{a}}(\bar{a}') = a'_2$  for  $\bar{a}'$  a permutation of  $\bar{a}$ )
  - (ii) if  $\bar{b} \in {}^3X$  is not one-to-one then  $f_{\bar{a}}(\bar{b}) = b_1$
- (b) for every one-to-one  $\bar{a} \in {}^3X$ , for some  $f = f_{\bar{a}} \in \mathcal{F}_{[3]}$  we have:
  - (i) if  $\bar{b}$  is a permutation of  $\bar{a}$  then  $f_{\bar{a}}(\bar{b}) = b_1$
  - (ii) if  $\bar{b} \in {}^3X$  is not one-to-one then  $f_{\bar{a}}(\bar{b}) = g_{r;1,2}(\bar{b})$
- (c) for every one-to-one  $\bar{a} \in {}^3X$ , for some  $f = f_{\bar{a}} \in \mathcal{F}_{[3]}$  we have
  - (i) if  $\bar{b}$  is a permutation of  $\bar{a}$  then  $f_{\bar{a}}(\bar{b}) = a_1$
  - (ii) if  $\bar{b} \in {}^3X$  is not one-to-one then  $f_{\bar{a}}(\bar{b}) = g_{r;1,2}(\bar{b})$

*Proof.* As  $\mathcal{F}$  is symmetric, it suffices to prove “for some  $\bar{a}$ ” instead of “for every  $\bar{a}$ ”.

Case 1: For some  $\ell(*)$  if  $\bar{b} \in {}^3X$  is not one to one then  $f^*(\bar{b}) = b_{\ell(*)}$ .

As  $f^*$  is not monarchical for some one-to-one  $\bar{a} \in {}^3X$ ,  $f^*(\bar{a}) \neq a_{\ell(*)}$  say  $f^*(\bar{a}) = a_{k(*)}$ ,  $k(*) \neq \ell(*)$ . As  $\mathcal{F}$  is symmetrical without loss of generality  $\ell(*) = 1$ ,  $k(*) = 2$ . So possibility (a) holds.

Case 2: Not case 1.

By 2.8, without loss of generality  $f^*$  satisfies (a),(b) or (c) of 2.8 with  $f^*$  instead of  $g$ . But clause (a) of 2.8 is case 1 above. So we can assume that case (b) of 2.8 holds, i.e.

(\*) if  $\bar{b} \in {}^3X$  is not one-to-one then  $f^*(\bar{b}) = g_{r;1,2}$ , i.e.

$$f^*(\bar{b}) = \begin{cases} b_2 & \text{if } b_2 = b_3 \\ b_1 & \text{if } b_2 \neq b_3 \end{cases}$$

If 2.10 applies we are done as (a) or (b) of 2.10 holds hence (a) or (b) of 3.1 respectively holds, so assume 2.10 does not apply. So consider a one-to-one sequence  $\bar{a} \in {}^3X$  and it follows that we have sequences  $\bar{a}^1, \bar{a}^2$ , both a permutation of  $\bar{a}$  such that  $\bigwedge_i [(f^*(\bar{a}^1) = a_i^1) \equiv (f^*(\bar{a}^2) \neq a_i^2)]$ .

Using closure under composition of  $\mathcal{F}$  and its being symmetric for every permutation  $\sigma$  of  $\{1, 2, 3\}$  (and as  $g_{r;1,2}$  is preserved by permuting the variables) there is  $f_\sigma \in \mathcal{F}_{[3]}$  such that:

- (i)  $f_\sigma(a_{\sigma(1)}, a_{\sigma(2)}, a_{\sigma(3)}) = a_1$
- (ii) if  $\bar{b} \in {}^3X$  not one-to-one then  $f(\bar{b}) = g_{r;1,2}(\bar{b})$ .

Let  $\langle \sigma_\rho : \rho \in {}^32 \rangle$  list the permutations of  $\{1, 2, 3\}$ , necessarily with repetitions. Now we define by downward induction of  $k \leq 3$ ,  $f_\rho \in \mathcal{F}$  for  $\rho \in {}^k2$  (sequences of zeroes and ones of length  $k$ ) as follows:

$$\ell g(\rho) = 3 \Rightarrow f_\rho = f_{\sigma_\rho}$$

$$\ell g(\rho) < 3 \Rightarrow f_\rho(x_1, x_2, x_3) = f_\rho(x_1, f_{\rho^\frown \langle 0 \rangle}(x_1, x_2, x_3), f_{\rho^\frown \langle 1 \rangle}(x_1, x_2, x_3)),$$

Easily (by downward induction):

- (\*)<sub>1</sub> if  $\bar{b} \in {}^3X$  is with repetitions then  $f_\rho(\bar{b}) = g_{r;1,2}(\bar{b})$  (as  $g_{r;1,2}$  act as majority).  
Now we prove by downward induction on  $k \leq 3$
- (\*)<sub>2</sub> if  $\bar{b}$  is a permutation of  $\bar{a}$ ,  $\rho \in {}^k2$ ,  $\rho \triangleleft \nu \in {}^32$  and  $f_\nu(\bar{b}) = a_1$  then  $f_\rho(\bar{b}) = a_1$ .

This is straight and so  $f_{< >}$  is as required in clause (c).  $\square_{3.1}$

Similarly

**3.2 Claim.** 1) If  $g_{r;\ell,k} \in \mathcal{F}$  then

- (a)  $g_{r;\ell_1,k_1} \in \mathcal{F}$  when  $\ell_1 \neq k_1 \in \{1, \dots, r\}$
- (b)  $g_{s;1,2} \in \mathcal{F}$  for  $s \geq r$  [seemingly not used].

*Proof.* 1)(a) Trivial.

(b) By induction on  $s$  so without loss of generality  $s = r + 1$ . Let  $g(x_1, \dots, x_{r+1}) = g_{r;1,2}(x_1, g_{r;1,2}(x_1, \dots, x_r), x_3, \dots, x_{r+1})$  so  $g \in \mathfrak{C}$  and

$$g(a, b, b, \dots, b) = g_{r;1,2}(a, g_{r;1,2}(a, b, \dots, b), b, \dots, b) = g_{r;1,2}(a, b, b, \dots) = b.$$

Now if  $g_{r+1;1,2}(a_1, \dots, a_{r+1}) = a_1$  then

Case 1:  $g_{r;1,2}(a_1, a_2, \dots, a_r) = a_1$  - trivial.

Case 2: Not 1; so  $(a_1, \dots, a_{r+1}) = (a, b, \dots, b, c)$  where  $b \neq c$  so easy to check.



§3 GETTING  $\mathfrak{C}$  IS FULL

3.1 *Lemma.* : Assume

- (a)  $r \geq 3$ ,  $\mathcal{F}$  is a clone on  $X$
- (\*)  $f_{r;1,2} \in \mathcal{F}$  or just
- (\*)<sup>-</sup> if  $\bar{a} \in {}^r X$  one-to-one for some  $f = f_{\bar{a}} \in \mathcal{F}$ ,  $f_{\bar{a}}(\bar{a}) = a_2$  and  $[\bar{b} \in {}^r X$   
not one-to-one  $\Rightarrow f_{\bar{a}}(\bar{b}) = b_1]$
- (b)  $\mathfrak{C}$  is a (non empty) family of choice functions for  $\binom{X}{k} = \{Y \subseteq X : |Y| = k\}$
- (c)  $\mathfrak{C}$  is closed under every  $f \in \mathcal{F}$
- (d)  $\mathfrak{C}$  is symmetric
- (e)  $k \geq r > 2, k \geq 5, |X| - k \geq 7, r$ .

Then  $\mathfrak{C}$  is full (i.e. every choice function is in).

*Proof.* Without loss of generality  $r \geq 4$  (if  $r = 3$  then clause (e) is fine also for  $r = 4$ , the case (\*) holds is O.K. by 2.7, and if (\*)<sup>-</sup> we repeat the proof of 2.7 for the case  $r = 3$ , only  $g(x_1, x_2, x_3, x_4) = f_{\langle a_1, a_2, a_3 \rangle}(x_1, \tau_2, \tau_3)$  where  $\tau_2 = f_{\langle a_1, a_2, a_4 \rangle}(x_1, x_2, x_4)$ ,  $\tau_3 = f_{\langle a_1, a_3, a_4 \rangle}(x_1, x_3, x_4)$  where for one-to-one  $\bar{a} \in {}^3 X$ ,  $f_{\bar{a}}$  is defined by the symmetry; this is the proof of 4.7). Assume

$$\boxtimes c_1^* \in \mathfrak{C}, Y^* \in \binom{X}{k}, c_1^*(Y^*) = a_1^* \text{ and } a_2^* \in Y^* \setminus \{a_1^*\}.$$

Question: Is there  $c \in \mathfrak{C}$  such that  $c(Y^*) = a_2^*$  and  $(\forall Y \in \binom{X}{k})(Y \neq Y^* \Rightarrow c(Y) = c_1^*(Y))$ ?

Choose  $c_2^* \in \mathfrak{C}$  such that

- (a)  $c_2^*(Y^*) = a_2^*$
- (b)  $n(c_2^*) = |\{Y \in \binom{X}{k} : c_2^*(Y) = c_1^*(Y)\}|$  is maximal under (a).

Easily  $\mathfrak{C}$  is not a singleton so  $n(c_2^*)$  is well defined.

3.2 Subfact: A positive answer to the question implies that  $\mathfrak{C}$  is full.

[Why? Easy.]

Hence if  $n(c_2^*) = \binom{|X|}{k} - 1$  we are done so assume not and let  $Z \in \binom{X}{k}, Z \neq Y^*, c_1^*(Z) \neq c_2^*(Z)$ .

Case 1: For some  $Z$  as above and  $c_3^* \in \mathfrak{C}$  we have

$$c_3^*(Y^*) \notin \{a_1^*, a_2^*\}$$

$$c_3^*(Z) \in \{c_1^*(Z), c_2^*(Z)\}.$$

If so, let  $a_3^* = c_3^*(Y^*)$  and  $a_4^* \in Y^* \setminus \{a_1^*, a_2^*, a_3^*\}$ , etc. so  $\langle a_1^*, \dots, a_r^* \rangle$  is one-to-one,  $a_\ell^* \in Y^*$ .

Let  $c_\ell^* \in \mathfrak{C}$  for  $\ell = 4, \dots$  be such that  $c_\ell^*(Y^*) = a_\ell^*$  exists as  $\mathfrak{C}$  is symmetric.

By assumption (a) we can choose  $f \in \mathcal{F}_{[r]}$  such that

$$\begin{cases} f(a_1^*, \dots, a_r^*) = a_2^*, \\ \bar{a} \in {}^r X \text{ is with repetitions} \rightarrow f(\bar{a}) = a_1 \end{cases}$$

Let  $c = f(c_1^*, c_2^*, \dots, c_r^*)$  so  $c \in \mathfrak{C}$  and:

$$c(Y^*) = f(a_1^*, a_2^*, \dots, a_r^*) = a_2^*$$

$$\begin{aligned} Y \in \binom{X}{k} \quad \& \quad c_1^*(Y) = c_2^*(Y) \Rightarrow c(Y) = f(c_1^*(Y), c_2^*(Y), \dots) \\ &= f(c_1^*(Y), c_1^*(Y), \dots) = c_1^*(Y) \end{aligned}$$

$$\begin{aligned} c(Z) &= f(c_1^*(Z), c_2^*(Z), c_3^*(Z), \dots) \\ &= c_1^*(Z) \text{ (as } |\{c_1^*(Z), c_2^*(Z), c_3^*(Z)\}| \leq 2). \end{aligned}$$

So  $c$  contradicts the choice of  $c_2^*$ .

Case 2: There are  $c_3^*, c_4^* \in \mathfrak{C}$  such that  $c_3^*(Y^*) \neq c_4^*(Y^*)$  are  $\neq a_1^*, a_2^*$  but  $c_3^*(Z) = c_4^*(Z)$  or at least  $|\{c_1^*(Z), c_2^*(Z), c_3^*(Z), c_4^*(Z)\}| < 4$ .

*Proof.* Similar.

Case 3: Not case 1 nor 2.

Let  $\mathcal{P} = \{Z : Z \subseteq X, |Z| = k \text{ and } c_1^*(Z) \neq c_2^*(Z)\}$  so

(\*)<sub>1</sub>  $\mathcal{P} \neq \emptyset, \{Y^*\}$ .

[Why?  $\mathcal{P} \neq \{Y^*\}$  by the subfact above. Also we can find  $Z \in \binom{X}{k}$  such that  $|Y^* \cap Z| = 2, c_1^*(Y^*) \notin Z$ . Let  $\pi \in \text{Per}(X)$  be the identity  $Z, \pi(c_1^*(Y^*)) \neq c_1^*(Y^*)$ . So conjugating  $c_1^*$  by  $\pi$  we get  $c_2^*, n(c_2^*) > 0$ .]

(\*)<sub>2</sub> if  $Z \in \mathcal{P}, c \in \mathfrak{C}$  and  $c(Z) \in \{c_1^*(Z), c_2^*(Z)\}$  then  $c(Y^*) \in \{c_1^*(Y^*), c_2^*(Y^*)\}$ .  
[Why? By not case 1.]

Subcase 3a: For some  $Z$  we have

$$\begin{cases} Z \in \mathcal{P} & \text{and :} \\ |Y^* \setminus Z| \geq 4 & \text{or just } |Y^* \setminus Z \setminus \{a_1^*, a_2^*\}| \geq 2 \text{ and } |Y^* \setminus Z| \geq 3 \end{cases}$$

Let  $b_1, b_2, b_3 \in Y^* \setminus Z$  be pairwise distinct. As  $\mathfrak{C}$  is symmetric there are  $d_1, d_2, d_3 \in \mathfrak{C}$  such that  $d_\ell(Y^*) = b_\ell$  for  $\ell = 1, 2, 3$ . The number of possible truth values of  $d_\ell(Z) \in Y^*$  is 2 so without loss of generality  $d_1(Z) \in Y^* \leftrightarrow d_2(Z) \in Y^*$  and we can forget  $b_3, d_3$ .

So for some  $\pi \in \text{Per}(X)$  we have  $\pi(Y^*) = Y^*, \pi(Z) = Z, \pi \upharpoonright (Y^* \setminus Z) = \text{identity}$  hence  $\pi(b_\ell) = b_\ell$  for  $\ell = 1, 2$  and  $\pi(d_1(Z)) = d_2(Z)$ , note that  $d_\ell(Z) \in Z$ , so this is possible; so without loss of generality  $d_1(Z) = d_2(Z)$ .

As  $|Y^* \setminus Z \setminus \{a_2^*, a_2^*\}| \geq 2$ , using another  $\pi \in \text{Per}(X)$  without loss of generality  $\{b_1, b_2\} \cap \{a_1^*, a_2^*\} = \emptyset$ . So  $d_1, d_2$  gives a contradiction by case 2.

*Remark.* This is enough for non polynomial  $|\mathfrak{C}|$  as  $|\{Y : |Y \setminus Z^*| \leq 3\}| \leq |Y|^6$ .

Case 3b: Not case 3a.

So  $Z \in \mathcal{P} \setminus \{Y^*\} \Rightarrow |Z \setminus Y^*| \leq 3$  hence (recalling  $|Z \setminus Y^*| = |Y^* \setminus Z|$ ) we have  $Z \in \mathcal{P} \setminus \{Y^*\} \Rightarrow |Z \cap Y^*| \geq k - 3 \geq 1$ .

Now

$\boxtimes_0$  for  $Z \in \mathcal{P} \setminus \{Y^*\}$  there is  $c^* \in \mathfrak{C}$  such that  $c^*(Y^*) \neq c^*(Z)$   
[Why? Otherwise “by  $\mathfrak{C}$  is symmetric” for any  $Z \in \mathcal{P} \setminus \{Y^*\}$  we have:

$$\circledast \quad c \in \mathfrak{C} \wedge Y', Y'' \in \binom{X}{k} \wedge |Y' \cap Y''| = |Z \cap Y^*| \Rightarrow c(Y') = c(Y'')$$

Define a graph  $\mathfrak{G} = \mathfrak{G}_Z$ : the set of nodes  $\binom{X}{k}$

the set of edge  $\{(Y', Y'') : |Y' \cap Y''| = |Y^* \cap Z|\}$

now this graph is connected: if  $\mathcal{P}_1, \mathcal{P}_2$  are nonempty disjoint set of nodes with union  $\binom{X}{k}$ , then there is a cross edge by 3.5 below (why? clause  $(\alpha)$  there is impossible by  $(*)_1$  and clause  $(\beta)$  is impossible by the first line of case 3). This gives contradiction to  $\circledast$ . So  $\boxtimes_0$  holds.]

We claim:

⊠<sub>1</sub> for  $Z \in \mathcal{P}$  and  $d \in \mathfrak{C}$  we have

$$d(Y^*) \in Z \cap Y^* \Rightarrow d(Z) = d(Y^*).$$

[Why? Assume  $d, Z$  forms a counterexample; as  $|Y^* \setminus Z| \leq 3$  and  $k \geq 7$  (see 3.1(e)) so if  $k \geq 8$  then  $|Y^* \cap Z| \geq k - 3 \geq 5$  so  $Y^* \cap Z \setminus \{a_1^*, a_2^*\}$  has  $\geq 3$  members looking again at subcase 3 this always holds. Now for some  $\pi_1, \pi_2 \in \text{Per}(X)$  we have  $\pi_1(Y^*) = Y^* = \pi_2(Y^*)$ ,  $\pi_1(Z) = Z = \pi_2(Z)$ ,  $\pi_1(d(Z)) = \pi_2(d(Z))$ ,  $\pi_1(d(Y^*)) \neq \pi_2(d(Y^*))$  are from  $Z \cap Y^* \setminus \{a_1^*, a_2^*\}$ ; recall we are assuming that  $d(Y^*) \in Z \cap Y^*$  and  $d(Z) \neq d(Y^*)$ . Let  $d_1, d_2$  be gotten from  $d$  by conjugating by  $\pi_1, \pi_2$ , so we get Case 2, contradiction.]

⊠<sub>2</sub> if  $d \in \mathfrak{C}, Y \in \binom{X}{k}$  and  $d(Y) = a$  then

$$(\forall Y')(a \in Y' \in \binom{X}{k} \rightarrow d(Y') = a).$$

[Why? By ⊠<sub>1</sub> + “ $\mathfrak{C}$  closed under permutations of  $X$ ”, we get: if  $k^* \in \{|Z \cap Y^*| : Z \in \mathcal{P} \setminus \{Y^*\}\}$  (which is not empty) then: if  $Z_1, Z_2 \in \binom{X}{k^*}$ ,  $|Z_1 \cap Z_2| = k^*$ ,  $d \in \mathfrak{C}$  and  $d(Z_1) \in Z_2$  then  $d(Z_1) = d(Z_2)$ . Clearly  $k^* < k$  (by  $Z \neq Y^*$ ) and  $Zk - k^* \leq |X|$ . First assume that we can choose such  $k^* > 0$ . Now for any  $d \in \mathfrak{C}$  and  $a \in X$ , claim 3.5 below applied to  $k^* - 1, k - 1, X \setminus \{a\}, (\{Y \setminus \{a\} : a \in Y \text{ and } d(Y) = a\}, \{Y \setminus \{a\} : a \in Y \text{ and } d(Y) \neq a\})$ . We can demand  $a \in \text{Rang}(d)$ . Now clause  $(\alpha)$  there gives the desired conclusion (for  $Y, a$  as in ⊠<sub>2</sub>). As we know  $k - k^* \leq 3, k \geq 7$  clause  $(\beta)$  is imposible too so we are done.]

Now we get a contradiction: as said above in ⊠<sub>0</sub> for some  $c^* \in \mathfrak{C}$  and  $Z \in \mathcal{P} \setminus \{Y^*\}$  we have  $c^*(Y^*) \neq c^*(Z)$ , choose  $Y \in \binom{X}{k}$  such that  $\{c^*(Y^*), c^*(Z)\} \subseteq Y$ . So by ⊠<sub>2</sub> we have  $d(Y) = d(Y^*)$  and also  $d(Y) = d(Z)$ , contradiction.  $\square_{3.1}$

**3.3 Claim.** In 3.1 we can replace (a) by

(a)<sup>\*</sup>(i)  $\mathcal{F}$  a clone on  $X$ ,

(ii)  $g^* \in \mathcal{F}_{[3]}$  where (note  $g^* = g_{3;1,2}$ )

$$g^*(x_1, x_2, x_3) = \begin{cases} x_2 & x_2 = x_3 \\ x_1 & \text{otherwise} \end{cases}$$

or just

(ii)<sup>-</sup> for  $\bar{a}^* \in {}^r X$  with no repetitions for some  $g = g_{\bar{a}^*}$ ,  $g(\bar{a}^*) = a_1^*$  and if  $\bar{a} \in {}^r X$  is with repetitions then  $g_{\bar{a}^*}(\bar{a}) = g^*(\bar{a})$ .

*Proof.* Let  $c_1^* \in \mathfrak{C}, Y^* \in \binom{X}{k}, a_1^* = c_1^*(Y^*), a_2^* \in Y^* \setminus \{a_1^*\}$ , we choose  $c_2^*$  as in the proof of 3.1.

Let  $\mathcal{P} = \{Y : Y \in \binom{X}{k}, Y \neq Y^*, c_1^*(Y) \neq c_2^*(Y)\}$ ; we assume  $\mathcal{P} \neq \emptyset$  and shall get a contradiction, (this suffices).

(\*)<sub>1</sub> there are no  $Z \in \mathcal{P}$  and  $d \in \mathfrak{C}$  such that

$$d(Y^*) = c_2^*(Y^*)$$

$$d(Z) \neq c_2^*(Z).$$

[Why? If so, let  $c = g(c_1^*, c_2^*, d)$ ,  $g$  is  $g^*$  or just any  $g_{\langle c_1^*(Z), c_2^*(Z), d(Z) \rangle}$  (from (a)\**iii*) of the assumption.

So  $c \in \mathfrak{C}$  and

- (A)  $c(Y^*) = g(c_1^*(Y^*), c_2^*(Y^*), d(Y^*)) = g(c_1^*(Y), c_2^*(Y^*), c_2^*(Y^*)) = c_2^*(Y^*)$
- (B)  $c(Z) = g(c_1^*(Z), c_2^*(Z), d(Z)) = c_1^*(Z)$  as  $d(Z) \neq c_2^*(Z)$   
(two cases: if  $\langle c_1^*(Z), c_2^*(Z), d(Z) \rangle$  with no repetitions - by the choice of  $g$ , otherwise it is equal to  $g^*(c_1^*(Z), c_2^*(Z), d(Z))$ )
- (C)  $Y \in \binom{Y}{k}, Y \neq Y^*, Z \notin \mathcal{P} \Rightarrow c_2^*(Y) = c_1^*(Y) \Rightarrow c(Y) = g(c_1^*(Y), c_2^*(Y), d(Y)) = g^*(c_1^*(Y), c_1^*(Y), d(Y)) = c_1^*(Y)$ .  $\square_{(*)_1}$

So (\*)<sub>1</sub> holds by  $c_2^*$ -s choice.]

(\*)<sub>2</sub> if  $\pi \in \text{Per}(X); \pi(Y^*) = Y^*, \pi(c_2^*(Y^*)) = c_2^*(Y^*)$  then

$$(\alpha) \quad Y \in \mathcal{P} \ \& \ \pi(Y) = Y \Rightarrow \pi(c_2^*(Y)) = c_2^*(Y)$$

$$(\beta) \quad Y \in \mathcal{P} \Rightarrow c_2^*(\pi(Y)) = \pi(c_2^*(Y)).$$

[Why? Otherwise may “conjugate”  $c_2^*$  by  $\pi^{-1}$  getting  $d \in \mathfrak{C}$  which gives a contradiction to (\*)<sub>1</sub>.]

(\*)<sub>3</sub> let  $Z \in \mathcal{P}$  then there are no  $d_1, d_2 \in \mathfrak{C}$  such that

$$d_1(Z) = d_2(Z) \neq c_2^*(Z)$$

$$d_1(Y^*) \neq d_2(Y^*).$$

[Why? Let  $g = g_{\langle c_2^*(Y^*), d_1(Y^*), d_2(Y^*) \rangle}$  be as in the proof of (\*)<sub>1</sub>. If the conclusion fails we let  $c = g(c_2^*, d_1, d_2)$  so

$$c(Y^*) = g(c_2^*(Y^*), d_1(Y^*), d_2(Y^*)) = c_2^*(Y^*) \text{ as } d_1(Y^*) \neq d_2(Y^*) + \text{choice of } g \text{ and } c(Z) = g(c_2^*(Z), d_1(Z), d_2(Z)) = d_1(Z) \neq c_2^*(Z) \text{ as } d_1(Z) = d_2(Z) \neq c_2^*(Z).$$

So  $c$  contradicts (\*)<sub>1</sub>.]

(\*)<sub>4</sub> for  $Z \in \mathcal{P}$ , there are no  $d_1, d_2 \in \mathfrak{C}$  such that  $d_1(Z) = d_2(Z), d_1(Y^*) \neq d_2(Y^*)$  except possibly when  $\{d_1(Z)\} = \{c_2^*(Z)\} \in \{Z \cap Y^*, Z \setminus Y^*\}$ .

[Why? If  $d_1(Z) \neq c_2^*(Z)$  use (\*)<sub>3</sub>, if  $d_1(Z) = c_2^*(Z)$  and there is  $\pi \in \text{Per}(X)$  satisfying  $\pi(Y^*) = Y^*, \pi(Z) = Z, \pi(c_2^*(Z)) \neq c_2^*(Z)$  we use it to get the situation in (\*)<sub>3</sub>.]

Let

$$K = \{(m) : \text{for some } Z \in \mathcal{P} \text{ we have } |Z \cap Y^*| = m\}$$

we are assuming  $K \neq \emptyset$ . By  $(*)_4$  symmetry we know

$$(*)_5 \text{ if } (m) \in K, 1 \neq m < k-1, c_1, c_2 \in \mathfrak{C} \text{ and } Z_1, Z_2 \in \binom{X}{k} \text{ satisfying } c_1(Z_1) = c_2(Z_1) \text{ and } |Z_1 \cap Z_2| = m, \underline{\text{then}} c_1(Z_2) = c_2(Z_2).$$

Case 1: There is  $(m) \in K$  such that  $1 \neq m < k-1$ , let  $\mathcal{P}' = \mathcal{P} \cup \{Y^*\}$ .

For any  $c_1, c_2 \in \mathfrak{C}$  let  $\mathcal{P}_{c_1, c_2} = \{Y \in \binom{Y}{k} : c_1(Y) = c_2(Y)\}$ .

By  $(*)_5$  we have  $[Y_1, Y_2 \in \binom{X}{k} \wedge |Y_1 \cap Y_2| = m \Rightarrow [Y_1 \in \mathcal{P}_{c_1, c_2} \equiv Y_2 \in \mathcal{P}_{c_1, c_2}]]$ .

Let  $Y_1 \in \binom{X}{k}, c_1 \in \mathfrak{C}$ , let  $a = c_1(Y_1)$  let  $Y_2 \in \binom{X}{k}$  be such that  $\{a, b\} = Y_1 \setminus Y_2$  for some  $b \neq a$ . By conjugation there is  $c_2 \in \mathfrak{C}$  such that  $c_2(Y_1) = a = c_1(Y_1)$  &  $c_1(Y_2) \neq c_2(Y_2)$ . To  $\mathcal{P}_{c_1, c_2}$  apply 3.5 below; so necessarily  $|X| = 2k, m = 0$ . But as  $m = 0, (m) \in K$  there is  $Y \in \mathcal{P}, |Y \cap Y^*| = m = 0$  hence  $Y = X \setminus Y^*$ , and by  $(*)_2(\alpha)$  we get a contradiction, i.e. we can find a  $\pi$  contradicting it.

Case 2:  $(m) \in K, m = k-1$  and not Case 1.

Let  $Z \in \mathcal{P}, |Z \cap Y^*| = k-1$  so by  $(*)_4$  and  $\mathfrak{C}$  being symmetric

$$(*)_6 \text{ if } Z_1, Z_2 \in \binom{X}{k}, |Z_1 \cap Z_2| = k-1, d_1, d_2 \in \mathfrak{C}, d_1(Z_1) = d_2(Z_1), d_1(Z_2) \neq d_2(Z_2) \text{ then } \{d_1(Z_1)\} = Z_1 \setminus Z_2.$$

Also

$$(*)_7 \text{ if } Z_1, Z_2 \in \binom{X}{k}, |Z_1 \cap Z_2| = k-1 \text{ then for no } d \in \mathfrak{C} \text{ do we have } d(Z_1) \neq d(Z_2) \text{ \& } \{d(Z_1), d(Z_2)\} \subseteq Z_1 \cap Z_2.$$

[Why? Applying  $\pi \in \text{Per}(X)$  we get a contradiction to  $(*)_6$ .]

This case (2) is finished by the following claim (and then we shall continue).

**3.4 Claim.** *Assume  $(a)^*(i)$  of 3.3 and  $(b), (c)$  of 3.1 and  $(*)_7$  above (on  $\mathfrak{C}$ ). Then  $\mathfrak{C}$  is full.*

*Proof.* Now

$$(*)_8 \text{ for every } Z_1, Z_2 \in \binom{X}{k}, |Z_1 \cap Z_2| = k-1 \text{ and } a \in Z_1 \cap Z_2 \text{ there is no } d \in \mathfrak{C} \text{ such that } d(Z_1) = d(Z_2) = a.$$

Why? Otherwise we can find  $Z_1, Z_2$  such that  $|Z_1 \cap Z_2| = k-1, d(Z_1) = d(Z_2) = a$  hence for every  $Z_1, Z_2 \in \binom{X}{k}$  such that  $|Z_1 \cap Z_2| = k-1$  and  $a \in Z_1 \cap Z_2$  there is such a  $d$  (using  $\pi \in \text{Per}(X)$ ).

Let  $Z_1, Z_2 \in \binom{X}{k}$  such that  $|Z_1 \cap Z_2| = k - 1$ . Let  $x \neq y \in Z_1 \cap Z_2$ . Choose  $d_1 \in \mathfrak{C}$  such that  $d_1(Z_1) = d_1(Z_2) = x$ .

Choose  $d_2 \in \mathfrak{C}$  such that  $d_2(Z_1) = d_2(Z_2) = y$ .

Choose  $d_3 \in \mathfrak{C}$  such that  $d_3(Z_1) = y, d_3(Z_2) \in Z_2 \setminus Z_1$ .

Why is it possible to choose  $d_3$ ? (Using  $\pi \in \text{Per}(X)$ ), otherwise (using  $(*)_7$ ) we have

⊗ if  $Y_1, Y_2 \in \binom{X}{k}, |Y_1 \cap Y_2| = k - 1$   
 $d \in \mathfrak{C}, d(Y_1) \in Y_1 \cap Y_2$  then  $d(Y_2) \in Y_1 \cap Y_2$   
 hence by  $(*)_7, d(Y_2) = d(Y_1)$   
 so for  $d \in \mathfrak{C}$  we have (by a chain of  $Y$ 's)

$$Y_1, Y_2 \in \binom{X}{k}, d(Y_1) \in Y_1 \cap Y_2 \Rightarrow d(Y_2) = d(Y_1).$$

Let  $c \in \mathfrak{C}, Y_1 \in \binom{X}{k}, x_1 = c(Y_1)$ . Let  $x_2 \in X \setminus Y_1, Y_2 = Y_1 \cup \{x_2\} \setminus \{x_1\}$ , so if  $c(Y_2) \in Y_1 \cap Y_2$  we get a contradiction, so  $d(Y_2) = x_2$ .

Let  $x_3 \in Y_1 \cap Y_2, Y_3 = Y_1 \cup Y_2 \setminus \{x_3\}$  so  $Y_3 \in \binom{X}{k}, |Y_3 \cap Y_1| = k - 1 = |Y_3 \cap Y_2|$  and clearly  $c(Y_1), c(Y_2) \in Y_3$ .

If  $c(Y_3) \notin Y_1$  then  $Y_3, Y_1$  contradict ⊗. If  $c(Y_3) \notin Y_2$  then  $Y_3, Y_2$  contradict ⊗.

But  $c(Y_3) \in Y_3 \subseteq Y_1 \cup Y_2$  contradiction. So  $d_3$  exists.

We shall use  $d_1, d_2, d_3, Z_1, Z_2$  to get a contradiction (thus proving  $(*)_8$ ).

Let  $\{z\} = Z_2 \setminus Z_1$  so  $\langle x, y, z \rangle$  is with no repetitions.

Let  $d = g(d_1, d_2, d_3)$  so with  $g = g^*$  or  $g = g_{\langle x, y, z \rangle}$

$$d(Z_1) = g(d_1(Z_1), d_2(Z_1), d_3(Z_1)) = g(x, y, y) = y$$

(see Definition of  $g$ )

$$\begin{aligned} d(Z_2) &= g(d_1(Z_2), d_2(Z_2), d_3(Z_2)) \\ &= g(x, y, z) = x \\ &\text{by Definition of } g \text{ as } y \neq z \text{ because } y \in Z_1, z \notin Z_1. \end{aligned}$$

So  $Z_1, Z_2, d$  contradicts  $(*)_7$ .

So we have proved  $(*)_8$ .

(\*)<sub>9</sub> if  $|Z_1 \cap Z_2| = k - 1, Z_1, Z_2 \in \binom{X}{k}, d \in \mathfrak{C}, d(Z_1) \in Z_1 \cap Z_2$ , then  $d(Z_2) \in Z_2 \setminus Z_1$ .  
 [Why? By  $(*)_7, d(Z_2) \notin Z_1 \cap Z_2 \setminus \{d(Z_1)\}$  and by  $(*)_8, d(Z_2) \notin \{d(Z_1)\}$ .]

Now we shall finish proving the claim 3.4.

Let  $c \in \mathfrak{C}$ .

Now let  $x_1, x_2 \in X$  be distinct and  $Y \subseteq X \setminus \{x_1, x_2\}$ ,  $|Y| = k$ . Let  $x_3 = c(Y)$ ,  $x_4 \in Y \setminus \{x_3\}$  and  $x_5 \in Y \setminus \{x_3, x_4\}$ .

So  $Y_1 = Y \cup \{x_1\} \setminus \{x_4\}$  belong to  $\binom{X}{k}$  satisfies  $|Y_1 \cap Y| = k-1$  and  $c(Y) = x_3 \in Y_1 \cap Y$  hence by  $(*)_9$ , we have  $c(Y_1) = x_1$ .

Let  $Y_2 = Y \cup \{x_2\} \setminus \{x_4\}$  so similarly  $c(Y_2) = x_2$ .

Let  $Y_3 = Y \cup \{x_1, x_2\} \setminus \{x_4, x_5\}$ , so  $Y_3 \in \binom{X}{k}$  and  $Y_3 \setminus Y_1 = \{x_2\}$  and  $Y_3 \setminus Y_2 = \{x_1\}$ .

If  $c(Y_3) \in Y$ , then

$$c(Y_3) \in Y_3 \cap Y = Y \setminus \{x_4, x_5\} \subseteq Y_1 \text{ hence } c(Y_3) \in Y_3 \cap Y_1$$

recall

$$c(Y_1) = x_1 \in Y_3 \cap Y_1$$

and  $c(Y_3) \neq x_1$  as  $x_1 \notin Y$  so  $(Y_3, Y_1, c)$  contradicts  $(*)_7$ .

If  $c(Y_3) = x_1$ , then  $c, Y_3, Y_1$  contradicts  $(*)_8$ .

If  $c(Y_3) = x_2$ , then  $c, Y_3, Y_2$  contradicts  $(*)_8$ .

Together contradiction, so we have finished proving 3.4 hence Case 2 in the proof of 3.1.  $\square_{3.4}$

Continuation of the proof of 3.1:

Case 3: Neither case 2 nor case 3.

As  $\mathcal{P} \neq \emptyset$  (otherwise we are done) clearly  $K = \{(1)\}$ . So easily (clearly  $2k-1 \leq |X|$  as  $(1) \in K$ ) and

$\boxtimes_1$  if  $|Y_1 \cap Y_2| = 1$ ,  $Y_1 \in \binom{X}{k}$ ,  $Y_2 \in \binom{X}{k}$  and  $d \in \mathfrak{C}$  then  $d(Y_1) \in Y_1 \cap Y_2 \vee d(Y_2) \in Y_1 \cap Y_2$ .

[Why? Otherwise by conjugation we can get a contradiction to  $(*)_4$  above.]

$\boxtimes_2$   $Y_1, Y_2 \in \binom{X}{k}$ ,  $|Y_1 \cap Y_2| = k-1$ ,  $d \in \mathfrak{C}$ ,  $d(Y_1), d(Y_2) \in Y_1 \cap Y_2$  is impossible.

[Why? Let  $x \in Y_1 \setminus Y_2$  and  $y \in Y_2 \setminus Y_1$ , we can find  $Y_3 \in \binom{X}{k}$  such that  $Y_3 \cap (Y_1 \cup Y_2) = \{x, y\}$  so  $Y_3 \cap Y_1 = \{x\}$ ,  $Y_3 \cap Y_2 = \{y\}$ ; this is possible as  $|X| \geq 2k-1$ . Apply  $\boxtimes_1$  to  $Y_3, Y_1, d$  and as  $d(Y_1) \neq x$  (as  $d(x_1) \in Y_Z$ ) we have  $c(Y_3) = x$ .

Apply  $\boxtimes_1$  to  $Y_3, Y_2, d$  and as  $d(Y_2) \neq y$  (as  $d(Y) \in Y_1$ ) we get  $d(Y_3) = y$ .

But  $x \neq y$ , contradiction.]

By  $\boxtimes_2$  we can use the proof of case 2 from  $(*)_7$ , i.e. Claim 3.4 to get contradiction.

$\square_{3.3}$



**3.5 Claim.** *Assume*

- (a)  $k^* < k < |X| < \aleph_0$ ,
- (b)  $\mathcal{P} \subseteq \binom{X}{k}$
- (c) if  $Z, Y \in \binom{X}{k}$ ,  $|Z \cap Y| = k^*$  then  $Z \in \mathcal{P} \Leftrightarrow Y \in \mathcal{P}$ .
- (d)  $2k - k^* \leq |X|$  (this is equivalent to clause (c) being non empty).

Then

- ( $\alpha$ )  $\mathcal{P} = \emptyset \vee \mathcal{P} = \binom{X}{k}$  or
- ( $\beta$ )  $|X| = 2k, k^* = 0$  and so  $E = E_{X,k} =: \{(Y_1, Y_2) : Y_1 \in \binom{X}{k}, Y_2 \in \binom{X}{k}, (Y_1 \cup Y_2 = X)\}$  is an equivalence relation on  $X$ , with each equivalence class is a doubleton and  $\mathcal{P}$  is a union of a set of  $E$ -equivalence classes.

*Proof.* If not clause ( $\alpha$ ), then for some  $Z_1 \in \mathcal{P}, Z_2 \in \binom{X}{k} \setminus \mathcal{P}$  we have  $|Z_1 \setminus Z_2| = 1$ .  
Let  $Z_1 \setminus Z_2 = \{a^*\}, Z_2 \setminus Z_1 = \{b^*\}$ .

Case 1:  $2k - k^* < |X|$ .

We can find a set  $Y^+ \subseteq X \setminus (Z_1 \cup Z_2)$  with  $k - k^*$  members (use  $|Z_1 \cup Z_2| = k + 1, |X \setminus (Z_1 \cup Z_2)| = |X| - (k + 1) \geq (2k - k^* + 1) - (k + 1) = k - k^*$ ).

Let  $Y^- \subseteq Z_1 \cap Z_2$  be such that  $|Y^-| = k^*$ .

Let  $Z = Y^- \cup Y^+$  so  $Z \in \binom{X}{k}, |Z \cap Z_1| = |Y^-| = k^*, |Z \cap Z_2| = |Y^-| = k^*$  hence  $Z_1 \in \mathcal{P} \Leftrightarrow Z \in \mathcal{P} \Leftrightarrow Z_2 \in \mathcal{P}$ , contradiction.

Case 2:  $2k - k^* = |X|$  and  $k^* > 0$ .

Let  $Y^+ = X \setminus (Z_1 \cup Z_2)$  so

$$|Y^+| = (2k - k^*) - |k + 1| = k - k^* - 1.$$

Let  $Y^- \subseteq Z_1 \cap Z_2$  be such that  $|Y^-| = k^* - 1$  (O.K. as  $|Z_1 \cap Z_2| = k - 1 \geq k^*$ ).

Let  $Z = Y^+ \cup Y^- \cup \{a^*, b^*\}$ . So  $|Z| = (k - k^* - 1) + (k^* - 1) + 2 = k, |Z_1 \cap Z| = |Y^- \cup \{a^*\}| = k^*, |Z_2 \cap Z| = |Y^- \cup \{b^*\}| = k^*$  and as in case 1 we are done.  $\square_{3.5}$

**3.6 Claim.** *If  $r(\mathcal{F}) < \infty$  then 3.1 or 3.3 apply so  $\mathfrak{C}$  is full.*

*Remark.* Recall  $r(\mathcal{F}) = \inf\{r : \text{some } f \in \mathcal{F}_{[r]} \text{ is not a monarchy}\}$ , see Definition 2.4.

*Proof.*

Case 1:  $r(\mathcal{F}) \geq 4$ .

Let  $f \in \mathcal{F}_{[r]}$  exemplify it, so by 2.5 for some  $\ell(*)$ :

$$\bar{a} \in {}^r X \text{ with repetitions} \Rightarrow f(\bar{a}) = a_{\ell(*)}.$$

As  $f$  is not a monarchy for some  $k(*) \in \{1, \dots, r\}$  and  $\bar{a}^* \in {}^r X$ ,  $f(\bar{a}^*) = a_{k(*)} \neq a_{\ell(*)}$ .

Without loss of generality  $\ell(*) = 1, k(*) = 2$  and 3.1 apply.

Case 2:  $r(\mathcal{F}) = 3$ .

Let  $f \in \mathcal{F}_{[r]}$  exemplify it. Now apply 3.1; if (a) there holds, apply 3.1, if (b) or (c) there holds, apply 3.3.

Case 3:  $r(\mathcal{F}) = 2$ .

By 4.7 below, clause (a) of 3.1 holds so we are done.

□<sub>3.6</sub>

§4 THE CASE  $r = 2$ 

This is revisited in §6 (non simple case), and we can make presentation simpler (e.g. 6.4).

## 4.1 Hypothesis.

- (a)  $r(\mathcal{F}) = 2$
- (b)  $|X| \geq 5$  (have not looked at 4).

**4.2 Claim.** Choose  $\bar{a}^* = \langle a_1^*, a_2^* \rangle, a_1^* \neq a_2^* \in X$ .

**4.3 Claim.** For some  $f \in \mathcal{F}$  and  $\bar{b} \in {}^2X$  we have

- (a)  $f(\bar{a}^*) = a_2^*$
- (b)  $\bar{a}^* \wedge \bar{b}$  is with no repetition
- (c)  $f(\bar{b}) = b_1 \neq b_2$ .

*Proof.* There is  $f \in \mathcal{F}$  not monarchical so for some  $\bar{b}, \bar{c} \in {}^2X$

$$f(\bar{b}) = b_1 \neq b_2, f(\bar{c}) = c_2 \neq c_1.$$

If  $\text{Rang}(\bar{b}) \cap \text{Rang}(\bar{c}) = \emptyset$  we can conjugate  $\bar{c}$  to  $\bar{a}^*$ ,  $f$  to  $f'$  which is as required.  
If not, find  $\bar{d} \in {}^2X, d_1 \neq d_2$  satisfying  $\text{Rang}(\bar{d}) \cap (\text{Rang}(\bar{a}) \cup \text{Rang}(\bar{b})) = \emptyset$  so  $\bar{d}, \bar{b}$  or  $\bar{d}, \bar{c}$  are like  $\bar{c}, \bar{b}$  or  $\bar{b}, \bar{c}$  respectively.  $\square_{4.3}$

**4.4 Claim.** There is  $f^* \in \mathcal{F}_{[2]}$  such that

- (a)  $f^*(\bar{a}^*) = \bar{a}_2^*$
- (b)  $b_1 \neq b_2 \in X, \{b_1, b_2\} \subseteq \{a_1^*, a_2^*\} \Rightarrow f(b_1, b_2) = b_2$
- (c)  $b_1 \neq b_2, \{b_1, b_2\} \not\subseteq \{a_1^*, a_2^*\} \Rightarrow f(b_1, b_2) = b_1$ .

*Proof.* Choose  $f$  such that

- (a)  $f \in \mathcal{F}_{[2]}$
- (b)  $f(\bar{a}^*) = a_2^*$
- (c)  $n(f) = |\{\bar{b} \in {}^2X : f(\bar{b}) = b_1\}|$  is maximal under (a) + (b).

Let  $\mathcal{P} = \{\bar{b} \in {}^2X : f(\bar{b}) = b_1\}$ . In each case we can assume that the previous cases do not hold for any  $f$  satisfying (a), (b), (c).

Case 1: There is  $\bar{b} \in {}^2(X \setminus \{a_1^*, a_2^*\})$  such that  $f(\bar{b}) = b_2 \neq b_1$ .

There is  $g \in \mathcal{F}_{[2]}$ ,  $g(\bar{a}^*) = a_2^*$ ,  $g(\bar{b}) = b_1$  (by 4.3 + conjugation). Let  $f^+(x, y) = f(x, g(x, y))$ .

So

$$(A) \quad f^+(\bar{a}^*) = f(a_1^*, g(\bar{a}^*)) = f(a_1^*, a_2^*) = a_2^*$$

$$(B) \quad f^+(\bar{b}) = f(b_1, g(\bar{b})) = f(b_1, b_1) = b_1$$

$$(C) \quad \text{if } \bar{c} \in \mathcal{P} \text{ then } f(\bar{c}) = c_1.$$

[Why does (C) hold? If  $g(\bar{c}) = c_1$  then  $f^+(\bar{c}) = f(c_1, g(\bar{c})) = f(c_1, c_1) = c_1$ .

If  $g(\bar{c}) = c_2$  then  $f^+(\bar{c}) = f(c_1, g(\bar{c})) = f(c_1, c_2) = f(\bar{c}) = c_1$  (last equality as  $\bar{c} \in \mathcal{P}$ ).]

By the choice of  $f$  the existence of  $f^+$  is impossible so

$$(*) \quad \bar{b} \in {}^2(X \setminus \{a_1^*, a_2^*\}) \Rightarrow f(\bar{b}) = b_1 \Rightarrow \bar{b} \in \mathcal{P} \text{ (if } b_1 = b_2 \text{ - trivial).}$$

Case 2: There are  $b_1 \neq b_2$  such that  $\{b_1, b_2\} \not\subseteq \{a_1^*, a_2^*\}$ ,  $f(b_1, b_2) = b_2$  and  $b_1 \neq a_1^* \wedge b_2 \neq a_2^*$ .

There is  $g \in \mathcal{F}_{[2]}$  such that  $g(a_1^*, a_2^*) = a_2^*$ ,  $g(b_1, b_2) = b_1$ .

[Why? There is  $\pi \in \text{Per}(X)$ ,  $\pi(b_1) = a_1^*$ ,  $\pi(b_2) = a_2^*$ ,  $\pi^{-1}(\{b_1, b_2\})$  is disjoint to  $\{a_1^*, a_2^*\}$ . Conjugate  $f$  by  $\pi^{-1}$ , getting  $g$  so  $g(a_1^*, a_2^*) = g(\pi b_1, \pi b_2) = \pi(f(b_1, b_2)) = \pi(b_2) = a_2^*$ ; let  $c_1, c_2$  be such that  $\pi(c_1) = b_1$ ,  $\pi(c_2) = b_2$  so

$$g(b_1, b_2) = g(\pi c_1, \pi c_2) = \pi(f(c_1, c_2)) = \pi(c_1) = b_1$$

(third equality as  $c_1, c_2 \notin \{a_2^*, a_2^*\}$  by not Case 1). So there is such  $g \in \mathcal{F}$ .]

Let  $f^+(x, y) = f(x, g(x, y))$ , as before  $f^+$  contradicts the choice of  $f$ .

Case 3: For some  $b' \neq b'' \in X \setminus \{a_1^*, a_2^*\}$  we have  $f(a_1^*, b') = b' \wedge f(a_1^*, b'') = a_1^*$ .

As in Case 2, using  $\pi \in \text{Per}(X)$  such that  $\pi(a_1^*) = a_1^*$ ,  $\pi(a_2^*) = a_2^*$ ,  $\pi(b') = b''$ .

Case 4: For some  $b' \neq b'' \in X \setminus \{a_1^*, a_2^*\}$  we have  $f(b', a_2^*) = a_2^* \wedge f(b'', a_2^*) = b''$ .

As in Case 3; recall that without loss of generality Case 1,2,3,4 fails.

Case 5: For some  $b', b'' \in X \setminus \{a_1^*, a_2^*\}$  we have  $f(a_1^*, b') = b' \wedge f(b'', a_2^*) = a_2^*$ .

As Cases 1,2,3,4 fail, without loss of generality  $b' \neq b''$  and prove as in Case 2 conjugating by  $\pi \in \text{Per}(X)$  such that  $\pi(b') = a_2^*$ ,  $\pi(a_1^*) = a_1^*$  and  $\pi(b'') = b''$  getting

$g$  which satisfies  $g(\bar{a}_1^*, a_2^*) = a_2^*$  and  $g(b', a_2^*) = \pi(f(b'', b')) = \pi(b'') = b'$  whereas  $f(b', a_2^*) = a_2^*$ ; so  $f^+(x, y) = f(x, g(x, y))$  contradicts the choice of  $f$ . Without loss of generality, Cases 1-5 fail.

Case 6: For some  $b \in X \setminus \{a_1^*, a_2^*\}$  we have  $f(a_1^*, b) = b$ .

So as Cases 1-5 fail we have

$$\circledast \quad \forall b_1, b_2 \in X [f(b_1, b_2) \neq b_1 \leftrightarrow (b_1 = a_1^* \ \& \ b_2 \neq a_1^*)]$$

(except in the case  $f(a_2^*, a_1^*) = a_1^*$  but then let  $\pi \in \text{Per}(X)$ ,  $\pi(a_1^*) = a_2^*$ ,  $\pi(a_2^*) = a_1^*$  (and  $\pi(a) = a$  for  $a \in X \setminus \{a_1^*, a_2^*\}$ ), then  $g = \pi f \pi^{-1}$  satisfies  $g(a_1^*, a_2^*) = a_2^*$  but for  $b \in X \setminus \{a_1^*, a_2^*\}$ ,  $g(a_1^*, b) = a_1^*$ , easy contradiction (or as below)).

Hence for every  $c \in X$  there is  $f_c \in \mathcal{F}_{[2]}$  such that

$$\circledast_{f_c} \quad \forall b_1 b_2 \in X [f_c(b_1, b_2) \neq b_1 \leftrightarrow (b_1 = c \ \& \ b_2 \neq c)].$$

Let  $a \neq c$  be from  $X$  and define  $f_{a,c} \in \mathcal{F}_{[2]}$  by  $f_{a,c}(x, y) = f_a(x, f_c(y, x))$ .

Assume  $b_1 \neq b_2$  so  $f_{a,c}^*(b_1, b_2) = b_2 \neq b_1$  implies  $f_c(b_2, b_1) \in \{b_1, b_2\}$ ,  $f_{a,c}(b_1, b_2) = f_a(b_1, f_c(b_2, b_1))$  and so (by the choice of  $f_a$ )  $b_1 = a$  &  $f_c(b_2, b_1) = b_2$  which (by the choice of  $f_c$ ) implies  $b_1 = a$  &  $b_2 \neq c$ . But  $b_1 = a$  &  $b_2 \neq c$  &  $b_1 \neq b_2$  implies  $f_c(b_2, b_1) = b_2$ ,  $f_{a,c}(b_1, b_2) = f_a(b_1, b_2) = b_2$ . So  $f_{a,c}(b_1, b_2) = b_2 \neq b_1 \leftrightarrow b_1 = a$  &  $b_2 \neq c$  &  $b_2 \neq b_1$ .

Let  $a = a_1^*$ . Let  $\langle c_i : i < i^* = |X| - 2 \rangle$  list  $X \setminus \{a_1^*, a_2^*\}$ . We define by induction on  $i \leq i^*$ , a function  $f_i \in \mathcal{F}_{[2]}$  by

$$f_0(x, y) = y$$

$$f_{i+1}(x, y) = f_i(x, f_{a, c_i}(x, y))$$

and let  $f' = f_{i^*}$ . Now by induction on  $i$  we can show  $f_i(a_1^*, a_2^*) = a_2^*$  and  $f'(b_1, b_2) = b_2 \neq b_1 \Rightarrow (\forall i < i^*)(f_{a, c_i}(b_1, b_2) = b_2 \neq b_1)$ .

So  $f' \in \mathcal{F}_{[2]}$ ,  $f'(a_1^*, a_2^*) = a_2^*$  and  $b_1 \neq b_2 \wedge (b_1, b_2) \neq (a_1^*, a_2^*), (a_2^*, a_1^*)$  implies  $f'(b_1, b_2) = b_1$ . By the choice of  $f$  (minimal  $n(f)$ ) we get a contradiction.

Case 7: For some  $b \in X \setminus \{a_1^*, a_2^*\}$  we have  $f(b, a_2^*) = a_2^*$ .

Similar to Case 6.

That is, as there, without loss of generality for every  $a \in X$  for some  $f_a \in \mathcal{F}_{[2]}$  we have

$$\circledast \quad (\forall b_1, b_2 \in X) [(f(b_1, b_2) = b_2 \neq b_1 \leftrightarrow b_2 = a \neq b_1)].$$

Let  $a \neq c \in X$  let  $f_{a,c}(x, y) = f_c(f_a(y, x), y)$ .

So for  $b_1 \neq b_2 \in X$

- (i)  $f_{a,c}(b_1, b_2) = b_2 (\neq b_1)$  implies  $f_a(f_c(b_2, b_1), b_2) = b_1$  which implies  $b_2 = c$  &  $f_c(b_2, b_1) = b_2$  which implies  $b_2 = c$  &  $b_1 \neq a$ .

We continue as there.

Case 8: Not Cases 1-7; not the conclusion.

So for  $\bar{a} = (a_1, a_2) = {}^2X$ ,  $a_1 \neq a_2$  there is  $f_{\bar{a}} \in \mathcal{F}$  such that

$$\{b_1, b_2\} \not\subseteq \{a_1, a_2\} \Rightarrow f_{\bar{a}}(b_1, b_2) = b_1$$

$$f_{\bar{a}}(a_1, a_2) = a_2$$

and (as “not the conclusion”)

$$f_{\bar{a}}(a_2, a_1) = a_2$$

Let  $\langle \bar{b}^i : i < i^* = |X|^2 - |X| - 2 \rangle$  list the pairs  $\bar{b} = (b_1, b_2) \in {}^2X$  such that  $b_1 \neq b_2$ ,  $\{b_1, b_2\} \neq \{a_1^*, a_2^*\}$ .

Define  $g_i \in \mathcal{F}_{[2]}$  by induction on  $i$ .

Let  $g_0(x, y) = x$ .

Let  $g_{i+1}(x, y) = f_{\bar{b}^i}(g_i(x, y), y)$ .

We can prove by induction on  $i \leq i^*$  that:  $g_i(a_1^*, a_2^*) = a_1^*$ ,  $g_i(a_2^*, a_1^*) = a_2^*$ ,  $j < i \Rightarrow g_i(\bar{b}^j) = b_2^j$ . So  $g_{i^*}$  is as required interchanging 1 and 2 that is  $g(x, y) =: g_{i^*}(y, x)$  is as required.  $\square_{4.4}$

**4.5 Definition/Choice.** For  $b \neq c \in X$  let  $f_{b,c}$  be like  $f$  in 4.4 with  $(b, c)$  instead of  $(a_1^*, a_2^*)$ , so  $f_{c,b}(c, b)$  is  $b$  and  $f(b, c) = c$  and  $f(x_{1,2}) = x_1$  if  $\{x_1, x_2\} \not\subseteq \{b, c\}$ .

**4.6 Claim.** Let  $a_1, a_2, a_3 \in X$  be pairwise distinct.

Then for some  $g \in \mathcal{F}_{[3]}$ :

- (i)  $\bar{b} \in {}^3X$  with repetitions  $\Rightarrow g(\bar{b}) = b_1$ ,  
(ii)  $g(a_1, a_2, a_3) = a_2$ .

*Proof.* Without loss of generality we replace  $a_2$  by  $a_3$  in (ii).

Let  $h_\ell$  for  $\ell = 1, 2, 3, 4$  be the three place functions

$$h_1(\bar{x}) = f_{a_1, a_2}(x_1, x_2)$$

$$h_2(\bar{x}) = f_{a_1, a_3}(x_1, x_3)$$

$$h_3(\bar{x}) = f_{a_2, a_3}(h_1(\bar{x}), h_2(\bar{x}))$$

$$h_4(\bar{x}) = f_{a_1, a_3}(x, h_3(\bar{x})).$$

Clearly  $h_1, h_2, h_3, h_4 \in \mathcal{F}_{[3]}$ . We shall show  $h_4$  is as required.

To prove clause (ii) note that for  $\bar{a} = (a_1, a_2, a_3)$  we have  $h_1(\bar{a}) = a_2, h_2(\bar{a}) = a_3, h_3(\bar{a}) = f_{a_2, a_3}(a_2, a_3) = a_3$  and  $h_4(\bar{a}) = f_{a_1, a_3}(a_1, a_3) = a_3$  as agreed above. To prove clause (i).

Let  $\bar{b} \in {}^3X$  be without repetitions such that  $\bar{b} \neq \bar{a}$ .

Case 1:  $b_1 \neq a_1, a_3$  so

$$h_4(\bar{b}) = f_{a_1, a_3}(b_1, h_3(\bar{b})) = b_1 \text{ as } b_1 \neq a_1, a_3.$$

Case 2:  $b_1 = a_1, b_2 \neq a_2$  hence  $b_1 \neq a_2, a_3$ , so

$$h_1(\bar{b}) = f_{a_1, a_2}(b_1, b_2) = b_1 \text{ (if } b_2 = a_1 \text{ also O.K.)}$$

$$h_3(\bar{b}) = f_{a_2, a_3}(h_1(\bar{b}), h_2(\bar{b})) = f_{a_2, a_3}(b_1, h_2(\bar{b})) = b_1 \text{ as } b_1 \neq a_2, a_3$$

$$h_4(\bar{b}) = f_{a_1, a_3}(b_1, h_3(\bar{b})) = h_{a_1, a_3}(b_1, b_1) = b_1.$$

Case 3:  $b_1 = a_1, b_2 = a_2, b_3 \neq a_3$ , so

$$h_1(\bar{b}) = f_{a_1, a_2}(b_1, b_2) = b_2$$

$$h_2(\bar{b}) = f_{a_1, a_3}(b_1, b_3) = f_{a_1, a_3}(a_1, b_3) = a_1 = b_1 \text{ as } b_3 \neq a_3 \text{ (if } b_3 = a_1 \text{ fine)}$$

$$h_3(\bar{b}) = f_{a_2, a_3}(h_1(\bar{b}), h_2(\bar{b})) = h_{a_2, a_3}(b_2, b_1) = b_2 \text{ as } b_1 = a_1 \neq a_2, a_3.$$

$$h_4(\bar{b}) = f_{a_1, a_3}(b_1, h_3(\bar{b})) = f_{a_1, a_3}(b_1, b_2) = b_1 \text{ as } b_2 = a_2 \neq a_1, a_3.$$

Case 4:  $b_1 = a_3, b_3 \neq a_1$ .  
So

$$h_1(\bar{b}) = f_{a_1, a_2}(b_1, b_2) = b_1 \text{ as } b_1 = a_3 \neq a_1, a_2$$

$$h_2(\bar{b}) = f_{a_1, a_3}(b_1, b_3) = f_{a_1, a_3}(a_3, b_3) = a_3 = b_1 \text{ as } b_3 \neq a_1$$

$$h_3(\bar{b}) = f_{a_2, a_3}(h_1(\bar{b}), h_2(\bar{b})) = f_{a_2, a_3}(b_1, b_1) = b_1$$

$$h_4(\bar{b}) = f_{a_1, a_3}(b_1, f_3(\bar{b})) = f_{a_1, a_3}(b_1, b_1) = b_1.$$

Case 5:  $b_1 = a_3, b_3 = a_1$ .

$$h_1(\bar{b}) = f_{a_1, a_2}(b_1, b_2) = b_1 \text{ as } b_1 = a_3 \neq a_1, a_2$$

$$h_2(\bar{b}) = f_{a_1, a_3}(b_1, b_3) = b_3 \text{ as } \{b_1, b_3\} = \{a_1, a_3\}$$

$$h_3(\bar{b}) = f_{a_2, a_3}(h_1(\bar{b}), h_2(\bar{b})) = f_{a_2, a_3}(b_1, b_3) \equiv b_1 \text{ as } b_3 = a_1 \neq a_2, a_3$$

$$h_4(\bar{b}) = f_{a_1, a_3}(b_1, f_3(\bar{b})) = f_{a_1, a_3}(b_1, b_1) = b_1.$$

as required. □<sub>4.6</sub>

**4.7 Claim.** *Let  $\bar{a}^* = (a_1^*, a_2^*, a_3^*, a_4^*) \in {}^4X$  be with no repetitions. Then for some  $g \in \mathcal{F}_{[4]}$  we have*

(i) *if  $\bar{b} \in {}^4X$  is with repetitions then  $f(\bar{b}) = b_1$*

(ii)  *$g(\bar{a}^*) = a_2^*$ .*



*Proof.* For any  $\bar{a} \in {}^3X$  with no repetitions let  $f_{\bar{a}}$  be as in 4.6 for the sequence  $\bar{a}$ . Let us define with  $(\bar{x} = (x_1, x_2, x_3, x_4)), g(\bar{x}) = g_0(x_1, g_2(x_1, x_2, x_4), g_3(x_1, x_3, x_4))$  with  $g_0 = f_{\langle a_1, a_2, a_3 \rangle}, g_2 = f_{\langle a_1, a_2, a_4 \rangle}, g_3 = f_{\langle a_1, a_3, a_4 \rangle}$ . So

- (A)  $g(\bar{a}^*) = g_0(a_1^*, g_2(a_1^*, a_2^*, a_3^*), g_3(a_1^*, a_3^*, a_4^*)) = g_0(a_1^*, a_2^*, a_3^*) = a_2^*$
- (B) if  $\bar{b} \in {}^4X$  and  $\langle b_1, b_2, b_4 \rangle$  is with repetition then  $g_2(b_1, b_2, b_4) = b_1$ , hence  $g(\bar{b}) = g_0(b_1, b_1, g_3(b_1, b_3, b_4)) = b_1$
- (C) if  $\bar{b} \in {}^4X$  and  $\langle b_1, b_3, b_4 \rangle$  is with repetition then  $g_3(a_1, a_3, a_4) = b_1$ , hence  $g(\bar{b}) = g_0(b_1, g_2(b_1, b_2, b_4), b_1) = b_1$
- (D)  $\bar{b} \in {}^4X$  is with repetitions, but neither (B) nor (C) then necessarily  $b_2 = b_3$  so  $\langle b_1, b_2, b_3 \rangle$  is with repetitions, so  $g(\bar{b}) = g_0(b_1, b_2, b_3) = b_1$ .

□?

→ scite{4.7} undefined

## PART B: NON SIMPLE CASE

## §5 FULLNESS FOR THE NON SIMPLE CASE

*5.1 Context.* As in §1:  $\mathfrak{C}$  is a  $(X, k)$ -FDF  $\mathcal{F} = \cup\{\mathcal{F}_{[r]} : r < \infty\}$  and  $\mathcal{F} = \{f : f \in \text{AV}(\mathfrak{C})\}$ , so

$\mathcal{F}_{[r]} = \{f : f \text{ is (not necessarily simple) function written}$

$f_Y(x_1, \dots, x_r), \text{ for } Y \in \binom{X}{k}, x_1, \dots, x_r \in Y \text{ such that}$

$f_Y(x_1, \dots, x_r) \in \{x_1, \dots, x_r\} \text{ and}$

$\mathfrak{C}$  is closed under  $f$ , i.e. if  $c_1, \dots, c_r \in \mathfrak{C}$

and  $c$  is defined by  $c = f(c_1, \dots, c_r)$

i.e.  $c(Y) = f_Y(c_1(Y), \dots, c_r(Y))$  then  $c \in \mathfrak{C}\}$

and we add (otherwise use Part A; alternatively combine the proofs).

*5.2 Hypothesis.* If  $f \in \mathcal{F}$  is simple then it is a monarchy.

**5.3 Definition.** 1)  $\mathcal{F}[Y] = \{f_Y : f \in \mathcal{F}\}$ .

2)  $\mathcal{F}_{[r]}(Y) = \{f_Y : f \in \mathcal{F}_{[r]}\}$ .

*5.4 Observation.* If  $f \in \mathcal{F}_{[r]}, Y \in \binom{X}{k}$ , then  $f_Y$  is an  $r$ -place function from  $Y$  to  $Y$  and

(\*)  $\mathcal{F}[Y]$  is a clone on  $Y$ .

**5.5 Definition.** 1)  $r(\mathcal{F}) = \text{Min}\{r : r \geq 2, \text{ some } f \in \mathcal{F}_{[r]} \text{ is not a monarchy}\}$  where

2)  $f$  is a monarchy iff for some  $t$  we have  $\forall Y \forall x_1, \dots, x_r \in Y [f_Y(x_1, \dots, x_r) = x_t]$ .

**5.6 Claim.** 1) For proving that  $\mathfrak{C}$  is full it is enough to prove, for some  $r \in \{3, \dots, k\}$

(\*) for every  $Y \in \binom{X}{k}$  and  $\bar{a} \in {}^r Y$  which is one to one there is  $f \in \mathcal{F}$  such that

(i)  $f_Y(\bar{a}) = a_2$

(ii) if  $Z \in \binom{X}{k}, Z \neq Y, \bar{b} \in {}^r Z$  then  $f_Z(\bar{b}) = b_1$ .

2) If  $r \geq 4$  we can weaken  $f_Z(\bar{b}) = b_1$  in clause (ii) to  $[b_3 = b_4 \vee b_1 = b_2 \vee b_1 = b_3 \vee b_2 = b_3] \rightarrow f_Y(\bar{b}) = b_1$  [not used].

*Proof.* The proof is as in the proof of ? below only we choose  $c_3, c_4, \dots, c_5$  such  
 $\rightarrow$  scite{5.8} undefined  
 that  $\bar{a} = \langle c_\ell(Y) : \ell = 1, 2, \dots, r \rangle$  is without repetitions and  $f = f_{\bar{a}}$  from (\*).  $\square_?$   
 $\rightarrow$  scite{5.6} undefined

**5.7 Claim.** In 5.6 we can replace (\*) by:  $r = 3$  and

- (\*) if  $Y \in \binom{X}{k}$  and  $\bar{a} \in {}^3Y$  one-to-one (or just  $a_2 \neq a_3$ ), then for some  $g \in \mathcal{F}_{[r]}$
- (i)  $g_Y(\bar{a}) = a_1$
  - (ii) if  $Z \in \binom{Y}{k}$ ,  $Z \neq Y$ ,  $\bar{b} \in {}^3Z$  is not one-to-one then  $g_Z(\bar{b}) = b$  if  $b_2 = b_3, b_1$  if otherwise (i.e.  $g_{3;1,2}(\bar{b})$ ).

*Proof.* Like 3.3. Let  $c_1^* \in \mathfrak{C}, Y^* \in \binom{X}{k}, a_1^* = c_2(Y^*), a_2^* \in Y^* \setminus \{a_1^*\}$  we choose  $c_2^*$  as in the proof of 5.6, i.e. 3.1.

Let  $\mathcal{P} = \{Y : Y \in \binom{X}{k}, Y \neq Y^*, c_1^*(Y) \neq c_2^*(Y)\}$ . As there it suffices to prove that  $\mathcal{P} = \emptyset$ . Now

$\boxtimes$  there are no  $Z \in \mathcal{P}$  and  $d \in \mathfrak{C}$  such that

$$d(Y^*) = c_2^*(Y^*)$$

$$d(Z) \neq c_2^*(Z).$$

[Why? If so, let  $c = g^*(c_1^*, c_2^*, d)$  where  $g$  is from (\*) for  $Z$ ,  $a_1 = c_1^*(Z)$ ,  $a_2 = c_2^*(Z)$ ,  $a_3 = d(Z)$ .]

Continue as there: the  $g_{\bar{a}}$  depends also on  $Y$ , and we write  $c(Y) = f_Y(c_1(Y), \dots, c_r(Y))$ .

$\square_{5.7}$

**5.8 Claim.** Assume  $r(\mathcal{F}) = 2$ ,  $(\mathfrak{C}, \mathcal{F})$  as usual) and

- (\*) for every  $a_1 \neq a_2 \in Y \in \binom{X}{k}$  for some  $f = f_{\langle a_1, a_2 \rangle}^Y \in \mathcal{F}$  we have
- (i)  $f_Y(\bar{a}) = a_2$
  - (ii)  $Z \in \binom{Y}{k}, Z \neq Y, \bar{b} \in {}^2Z \Rightarrow f_Z(\bar{b}) = b_1$ .

Then  $\mathfrak{C}$  is full.

*Remark.*  $\mathfrak{C}$  is full iff every choice function of  $\binom{X}{k}$  belongs to it.

*Proof.* If  $\mathfrak{C}$  is not full, as  $\mathfrak{C} \neq \emptyset$  there are  $c_1 \in \mathfrak{C}, c_0 \notin \mathfrak{C}, c_0$  a choice function for  $\binom{X}{k}$ . Choose such a pair  $(c_1, c_0)$  with  $|\mathcal{P}|$  minimal where  $\mathcal{P} = \{Y \in \binom{X}{k} : c_1(Y) \neq c_0(Y)\}$ . So clearly  $\mathcal{P}$  is a singleton, say  $\{Y\}$ . By symmetry for some  $c_2 \in \mathfrak{C}$  we have  $c_2(Y) = c_0(Y)$ . Let  $f$  be  $f_{c_1(Y), c_0(Y)}^Y = f_{c_1(Y), c_2(Y)}^Y$  from the assumption so  $f \in \mathcal{F}$  and let  $c = f(c_1, c_2)$  so clearly  $c \in \mathfrak{C}$  (as  $\mathfrak{C}$  is closed under every member of  $\mathcal{F}$ ).

Now

$$(A) \ c(Y) = f_Y(c_1(Y), c_2(Y)) = c_2(Y) = c_0(Y)$$

$$(B) \text{ if } Z \in \binom{X}{k} \setminus \{Y\} \text{ then}$$

$$c(Z) = f_Z(c_1(Z), c_2(Z)) = c_1(Z) = c_0(Y).$$

So  $c = c_0$  hence  $c_0 \in \mathfrak{C}$ , contradiction.  $\square_{5.8}$

**5.9 Claim.** Assume  $r(\mathcal{F}) = 2$  and  $\boxtimes(f^*)$  of 6.9 (see Definitions ?, 6.6) below  
 $\Rightarrow$  *scite*{6.3} undefined  
 holds. Then  $\mathfrak{C}$  is full.

*Proof.* We use conventions from 6.6, 6.7, 6.9 below. In  $\boxtimes(f^*)$  there are two possibilities.

Possibility (i):

This holds by 5.8.

Possibility (ii):

Similar to the proof of 5.8. Again  $\mathcal{P} = \{Y\}$  where  $\mathcal{P} = \{Y \in \binom{X}{k} : c_1(Y) \neq c_0(Y)\}$ . We choose  $c_2 \in \mathfrak{C}$  such that  $c_2(Y) = c_0(Y)$  &  $c_2(X \setminus Y) = c_0(Y)$ , continue as before. Why is this possible? Let  $\pi \in \text{Per}(X)$  be such that  $\pi(Y) = Y, \pi(c_1(Y)) = c_0(Y), \pi(c_1(X \setminus Y)) = c_1(X \setminus Y)$  (and of course  $\pi(X \setminus Y) = X \setminus Y$ ). Now conjugating  $c_1$  by  $\pi$  gives  $c_2$  as required.  $\square_{5.9}$

**5.10 Claim.** *If  $r(\mathcal{F}) < \infty$  then  $\mathfrak{C}$  is full.*

*Proof.* Let  $r = r(\mathcal{F})$ .

Case 1:  $r = 2$ .

So hypothesis 6.1 below holds.

If  $\boxtimes(f)$  of 6.9 holds for some  $f \in \mathcal{F}_{[r]}$ , by 5.9 we know that  $\mathfrak{C}$  is full. If  $\boxtimes(f)$  of 6.9 fails for every  $f \in \mathcal{F}_{[r]}$  then hypothesis 6.11 holds hence 6.12-6.18 holds. So by 6.18 we know that  $(*)$  of 5.6 holds (and  $\mathcal{P}_{\pm}$  is a singleton). So by 5.6,  $\mathfrak{C}$  is full.

Case 2:  $r \geq 4$ .

So hypothesis 7.1 holds. By 7.5 clearly  $(*)$  of 5.6 holds hence by 5.6 we know that  $\mathfrak{C}$  is full.

Case 3:  $r = 3$ .

Let  $f^* \in \mathcal{F}_{[3]}$  be not a monarchy. So for  $\bar{b} \in {}^3Y$  not one-to-one,  $Y \in \binom{X}{3}$ , clearly  $f_Y^*(\bar{b})$  does not depend on  $Y$ , so we write  $f^-(\bar{b})$ . If for some  $\ell(*)$ ,  $f^-(\bar{b}) = b_{\ell(*)}$  for every such  $\bar{b}$  then easily 5.6 apply. If  $f^-(\bar{b}) = g_{r;1,2}(\bar{b})$ , let  $\bar{a} \in {}^3Y$ ,  $Y \in \binom{X}{k}$ ,  $\bar{a}$  is one-to-one, so  $f_Y(\bar{b}) = a_k$  for some  $k$ ; by permuting the variables,  $f^-$  does not change while we have  $k = 1$ , so 5.7 apply. If both fail, then by repeating the proof of 2.8 of Part A, for some  $f' \in \mathcal{F}_{[3]}$ , for  $\bar{b}$  not one-to-one we have  $f_{\langle 1,2,1 \rangle}(\bar{b})$  or for  $\bar{b}$  not one to one  $f'_Y(\bar{b}) = f_{\langle 1,2,2 \rangle}(\bar{b})$ . By the last paragraph of the proof of ? we can

→ scite{2.8} undefined

assume the second case holds. In this case repeat the proof of the case  $\eta = \langle 1, 2, 2 \rangle$  in the end of the proof of ?.

→ scite{2.8} undefined

§6 THE CASE  $r(\mathcal{F}) = 2$ 

For this section

6.1 *Hypothesis.*  $r = 2$ .

6.2 Discussion: So  $(\alpha)$  or  $(\beta)$  holds where

$(\alpha)$  there are  $Y \in \binom{X}{k}$  and  $f \in \mathcal{F}_{[r]}(Y)$  which is not monarchy. Hence by §4, i.e. 4.4 for  $a \neq b \in Y$  there is  $f = f_{a,b}^Y \in \mathcal{F}_2[Y]$

$$f_Y(x, y) = \begin{cases} y & \text{if } \{x, y\} = \{a, b\} \\ x & \text{if otherwise} \end{cases}$$

$(\beta)$  every  $f_Y$  is a monarchy but some  $f \in \mathcal{F}_{[r]}$  is not.

6.3 **Definition/Choice.** Choose  $f^* \in \mathcal{F}_2$  such that

- (a)  $\neg(\forall Y)(\forall x, y \in Y)(f(x, y) = x)$
- (b) under (a),  $n(f) = |\text{dom}_1(f)|$  where  $\text{dom}_1(f) = \{(Z, a, b) : f_Z(a, b) = a \neq b \text{ and } Z \in \binom{X}{k} \text{ and } \{a, b\} \subseteq Z \text{ of course}\}$ , is maximal.

6.4 *Fact.* If  $f_1, f_2 \in \mathcal{F}_{[2]}$  and  $f$  is  $f(x, y) = f_1(x, f_2(x, y))$  (formally  $f(Y, x, y) = f_1(Y, x, f_2(Y, x, y))$  but we shall be careless) then  $\text{dom}_1(f) = \text{dom}_1(f_1) \cup \text{dom}_1(f_2)$ .

*Proof.* Easy.

6.5 **Claim.** If  $Z \in \binom{X}{k}$ ,  $f_Z^*(a^*, b^*) = b^* \neq a$  then

- (a)  $(\forall x, y \in Z)[f_Z^*(x, y) = y]$  or
- (b)  $x, y \in Z \wedge \{x, y\} \not\subseteq \{a^*, b^*\} \Rightarrow f_Z^*(x, y) = x$ .

*Proof.* As in 4.4 (+ 6.3 + 6.4), recalling that  $\mathcal{F}(Z)$  is a clone.

□<sub>6.5</sub>

**6.6 Definition.** Let

- (1)  $\mathcal{P}_1 = \mathcal{P}_1(f^*) = \{Z \in \binom{X}{k} : (\forall a, b \in Z)(f_Z^*(a, b) = a)\}$
- (2)  $\mathcal{P}_2 = \mathcal{P}_2(f^*) = \{Z \in \binom{X}{k} : (\forall a, b \in Z)(f_Z^*(a, b) = b)\}$
- (3)  $\mathcal{P}_\pm = \mathcal{P}_\pm(f^*) = \binom{X}{k} \setminus \mathcal{P}_1(f^*) \setminus \mathcal{P}_2(f^*)$ .

**6.7 Claim.** For  $Y \in \binom{X}{k}$  we have

- 1)  $Y \in \mathcal{P}_\pm(f^*)$  iff  $Y \in \binom{X}{k}$  and  $(\exists a, b \in Y)(f_Y^*(a, b) = a \neq b)$  and  $(\exists a, b \in Y)(f_Y^*(a, b) = b \neq a)$ .
- 2) If  $Y \in \mathcal{P}_\pm$ , then there are  $a_Y \neq b_Y \in Y$  such that  $f_Y^*(a_Y, b_Y) = b_Y$  and

$$\{a, b\} \subseteq Y \wedge \{a, b\} \not\subseteq \{a_Y, b_Y\} \Rightarrow f_Y^*(a, b) = a.$$

*Proof.* By 6.5.

**6.8 Claim.** 1)  $\langle \mathcal{P}_1, \mathcal{P}_2, \mathcal{P}_\pm \rangle$  is a partition of  $\binom{X}{k}$ .

2) For  $Y \in \mathcal{P}_\pm$  the pair  $(a_Y, b_Y)$  is well defined (but maybe  $(b_Y, a_Y)$  can serve as well).

*Proof.* By 6.5.

**6.9 Claim.** If  $\mathcal{P}_2(f^*) \neq \emptyset$  then

$\boxtimes_{f^*}(i)$   $\mathcal{P}_2 = \mathcal{P}_2(f^*)$  is a singleton,  $\mathcal{P}_\pm = \emptyset$  or

(ii)  $2k = |X|$ ,  $\mathcal{P}_2$  is a  $\{Y^*, Y^{**}\} \subseteq \binom{X}{k}$  where  $Y^* \cup Y^{**} = X$  and  $\mathcal{P}_\pm = \emptyset$ .

*Proof.* Assume  $\mathcal{P}_2 \neq \emptyset$ , let  $Y^* \in \mathcal{P}_2$ . As  $f^*$  is not a monarchy

$$(*)_1 \quad \mathcal{P}_1 \cup \mathcal{P}_\pm \neq \emptyset.$$

By Definition 6.6 and Fact 6.4,  $f^* \in \mathcal{F}_{[r]}$  satisfies

$(*)_2(i)$   $f_{Y^*}^*(a, b) = b$  for  $a, b \in Y^*$

(ii) if  $g \in \mathcal{F}_{[r]}$ ,  $g_{Y^*}(a, b) = b$  for  $a, b \in Y^*$  then  $\text{dom}_1(f^*) \supseteq \text{dom}_1(g)$ .

Hence

- (\*)<sub>3</sub> if  $Y_1 \in \mathcal{P}_2, Y_1 \notin \mathcal{P}_2, k^* = |Y_1 \cap Y_2|$  and  $Y \in \binom{Y}{k}, |Y \cap Y^*| = k^*$ , then  
 $Y \notin \mathcal{P}_2$  (even  $Y \in \mathcal{P}_1 \leftrightarrow Y_2 \in \mathcal{P}_1$ ).  
 [Why? By (\*)<sub>2</sub> as we can conjugate  $f^*$  by  $\pi \in \text{Per}(X)$  which maps  $Y^*$  onto  $Y^*$  and  $Y_1$  to  $Y$ .]

So by 3.5 (applied to all  $k^*$ )

- (\*)<sub>4</sub>(i)  $\mathcal{P}_2$  is the singleton  $\{Y^*\}$  or  
 (ii)  $\mathcal{P}_2$  is a  $\{Y^*, Y^{**}\}, 2k = |X|$  and  $Y^{**} = X \setminus Y^*$

- (\*)<sub>5</sub> if  $Z \in \mathcal{P}_\pm$ , then  $(\alpha)$  or  $(\beta)$

$$(\alpha) \quad \{a_Z, b_Z\} = Z \cap Y^*, f_Z^*(b_Z, a_Z) = a_Z$$

$$(\beta) \quad \{a_Z, b_Z\} = Z \setminus Y^*, f_Z(b_Z, a_Z) = a_Z.$$

[Why? Otherwise as  $k \geq 3$  we can choose  $\pi \in \text{Per}(X), \pi(Y^*) = Y^*, \pi(Z) = Z$  such that  $\pi''\{a_Z, b_Z\} \not\subseteq \{a_Z, b_Z\}$  and use 6.3, 6.4 on a conjugate of  $f^*$ .]

It is enough by (\*)<sub>4</sub> to prove  $\mathcal{P}_\pm = \emptyset$ .

So assume toward contradiction  $\mathcal{P}_\pm \neq \emptyset$ .

By (\*)<sub>5</sub> one of the following two cases occurs.

Case 1:  $Z^* \in \mathcal{P}_\pm, |Z^* \cap Y^*| = k - 2$ .

As we are allowed to assume  $k+2 < |X|$  there is  $Y \in \binom{X}{k}$  such that  $|Y \cap Y^*| = k-1$  and  $Y \cap Z^* = Y^* \cap Z^*$ . Now (by (\*)<sub>5</sub>) we have  $Y \notin \mathcal{P}_\pm$  and (by (\*)<sub>4</sub>) we have  $Y \notin \mathcal{P}_2$  so  $Y \in \mathcal{P}_1$ . So there is  $\pi \in \text{Per}(X)$  such that  $\pi(Y^*) = Y, \pi \upharpoonright Z^* = \text{identity}$ , let  $f = (f^*)^\pi$  so by 6.4 we get a contradiction to the choice of  $f^*$ .

Case 2:  $Z^* \in \mathcal{P}_\pm, |Z^* \cap Y^*| = 2$ .

A proof similar to case 1 works if  $Z^* \cup Y^* \neq X$ . So assume  $Z^* \cup Y^* = X$  (hence  $2k - 2 = |X|$ ).

Let  $\pi \in \text{Per}(X)$  be the identity on  $Z^* \cap Y^*$  and interchange  $Z^*, Y^*$ . Apply 6.4 on  $f^*, (f^*)^\pi$  so  $(a_{Z^*}, b_{Z^*}) \in \text{dom}_1(f^*) \cap \text{dom}_1((f^*)^\pi)$ , etc. easy contradiction.

□<sub>6.9</sub>

*6.10 Remark.* if  $\boxtimes(f^*)$  of 6.9 holds for some  $f^*$  then  $\mathfrak{C}$  (in the context of §5)  $\mathfrak{C}$  is full by 5.9.

*6.11 Hypothesis.* For no  $f \in \mathcal{F}_{[r]}$  is  $\boxtimes(f)$ .

*6.12 Conclusion.* 1)  $\mathcal{P}_2(f^*) = \emptyset$ .

2)  $\mathcal{P}_\pm \neq \emptyset$ .



- 3)  $\mathcal{P}_1 \neq \emptyset$ .  
 4) If  $Y \in \mathcal{P}_\pm$  and  $|Y \cap Z_1| = |Y \cap Z_2|$  and  $a_Y \in Z_1 \leftrightarrow a_Y \in Z_2$  and  $b_Y \in Z_1 \leftrightarrow b_Y \in Z_2$  where, of course,  $Y, Z_1, Z_2 \in \binom{X}{k}$ , then  $Z_1 \in \mathcal{P}_\pm \leftrightarrow Z_2 \in \mathcal{P}_\pm$ .

*Proof.* 1) By 6.11 + 6.9.

2) Otherwise  $f^*$  is a monarchy.

3) Assume not, so  $\mathcal{P}_\pm = \binom{X}{k}$ . Let  $Y \in \mathcal{P}_\pm, Z \in \binom{X}{k}, Z \cap \{a_Y, b_Y\} = \emptyset$  and<sup>2</sup>  $|Z \cap Y| > 2$  and  $|Z \setminus Y| > 2$ , we can get a contradiction to  $n(f^*)$ -s minimality.

4) By 6.3 and 6.4 as we can find  $\pi \in \text{Per}(X)$  such that  $\pi(Y) = Y, \pi(Z_1) = Z_2, \pi(a_Y) = a_Y, \pi(b_Y) = b_Y$ .  $\square_{6.12}$

**6.13 Claim.** *If  $Y, Z \in \mathcal{P}_\pm$  and  $Y \neq Z$ , then there is no  $\pi \in \text{Per}(X)$  such that:*

$$\pi(Y) = Y, \pi(Z) = Z$$

$$\pi(a_Y) = a_Y, \pi(b_Y) = b_Y$$

$$\{\pi(a_Z), \pi(b_Z)\} \not\subseteq \{a_Z, b_Z\}.$$

*Proof.* By 6.3 + 6.4.

**6.14 Claim.** *If  $Y \in \mathcal{P}_\pm, Z \in \mathcal{P}_\pm, 2 < |Y \cap Z| < k - 2$  then  $\{a_Z, b_Z\} = \{a_Y, b_Y\}$ .*

*Proof.* By 6.13.

**6.15 Claim.** *If  $Z_0, Z_1 \in \mathcal{P}_\pm$  and  $|Z_1 \setminus Z_0| = 1$  then  $\{a_{Z_0}, b_{Z_0}\} = \{a_{Z_1}, b_{Z_1}\}$ .*

*Proof.* We shall choose by induction  $i = 0, 1, 2, 3, 4$  a set  $Z_i \in \mathcal{P}_\pm$  such that  $j < i \Rightarrow |Z_i \setminus Z_j| = i - j$ , this holds for  $i = 1, j = 0$ , also this is sufficient as by 6.14 we have  $i = j + 3 \leq 4 \Rightarrow \{a_{Z_i}, b_{Z_i}\} = \{a_{Z_j}, b_{Z_j}\}$  as this applies to  $(j, i) = (0, 4)$  and  $(j, i) = (1, 4)$  we get the desired conclusion by transitivity of equality.

To choose  $Z_i$ , let  $x_i \in X \setminus Z_0 \cup \dots \cup Z_{i-1}$ ; possible as we exclude  $k + i - 1$  elements and choose  $y_i \in Z_0 \cap \dots \cap Z_{i-1} \setminus \{a_{Z_{i-1}}, b_{Z_{i-1}}\}$ . Now let  $Z_i = Z_{i-1} \cup \{y_i\} \setminus \{x_i\}$  easily  $j < i \Rightarrow |Z_i \setminus Z_j| = i - j$  and  $Z_i \in \mathcal{P}_\pm$  by 6.12(4) with  $Y, Z_1, Z_2$  there standing for  $Z_{i-1}, Z_{i-2}, Z_i$  here.

---

<sup>2</sup>I am sure that after checking we can save

6.16 Choice:  $Y^* \in \mathcal{P}_\pm$ .

*6.17 Conclusion.*

- (a)  $Y^* \in \mathcal{P}_\pm$
- (b) if  $Y \in \mathcal{P}_\pm$  then  $\{a_Y, b_Y\} = \{a_{Y^*}, b_{Y^*}\}$
- (c) one of the following possibilities holds
  - ( $\alpha$ )  $\mathcal{P}_\pm = \{Y^*\}$
  - ( $\beta$ )  $\mathcal{P}_\pm = \{Y \in \binom{X}{k} : \{a_{Y^*}, b_{Y^*}\} \subseteq Y\}$
  - ( $\gamma$ )  $\mathcal{P}_\pm = \{Y^*, Y^{**}\}$  where  $Y^{**} = (X \setminus Y^*) \cup \{a_{Y^*}, b_{Y^*}\}$  and  $|X| = 2k - 2$ .

*Proof of 6.17.* Note that

(\*) if  $Y_1, Y_2, Y_3 \in \mathcal{P}_\pm, |Y_1 \setminus Y_2| = 1$  and  $Y_3 \in \mathcal{P}_\pm, Y_4 \in \binom{X}{k}, |Y_3 \setminus Y_4| = 1$  and  $\{a_{Y_3}, b_{Y_3}\} = \{a_{Y_1}, b_{Y_1}\} \subseteq Y_4$  then  $Y_4 \in \mathcal{P}_\pm$  (hence  $\{a_{Y_4}, b_{Y_4}\} = \{a_{Y_3}, b_{Y_3}\} = \{a_{Y_1}, b_{Y_1}\}$ .  
 [Why? As there is a permutation  $\pi$  of  $X$  such that  $\pi(a_{Y_1}) = a_{Y_2}, \pi(b_{Y_1}) = Y_1, \pi(Y_3) = Y_1, \pi(Y_4) = Y_2$ . By ? we get a contradiction to the choice of  
 $\rightarrow$  scite{6.4} undefined  
 $f^*.$ ]

By the choice of  $Y^* \in \mathcal{P}_\pm$ , we have (a), now (b) follows from (c) by 6.15 by induction on  $|Y \setminus Y^*|$ , so it is enough to prove (c). Assume ( $\alpha$ ), ( $\gamma$ ) fail and we shall prove ( $\beta$ ). So there is  $Z_1 \in \mathcal{P}_\pm$  such that  $Z_1 \notin \{Y^*, (X \setminus Y^*) \cup \{a_{Y^*}, b_{Y^*}\}\}$ . We can find  $c_1, c_2 \in X \setminus \{a_{Y^*}, b_{Y^*}\}$  such that  $c_1 \in Y^* \leftrightarrow c_2 \in Y^*$  and  $c_1 \in Z_1 \leftrightarrow c_2 \notin Z_1$ . [Why? if  $Y^* \cup Z_1 \neq X$  any  $c_1 \in X \setminus Y^* \setminus Z_1, c_2 \in Z_1 \setminus Y^*$  will do; so assume  $Y^* \cup Z_1 = X$  so as  $k + 2 < |X|$  clearly by 6.14,  $|Z_1 \cap Y^*| \leq 2$ . As not case ( $\gamma$ ) of (c), that is by the choice of  $Z_1$ , necessarily  $\{a_{Y^*}, b_{Y^*}\} \not\subseteq Y^* \cap Z_1$  and using  $\pi \in \text{Per}(X), \pi \upharpoonright Z_1 = \text{id}, \pi(Y^*) = Y^*$  by 6.13 we contradict 6.3 + 6.4.]

Let  $Z_2 = Z_1 \cup \{c_1, c_2\} \setminus (Z_1 \cap \{c_1, c_2\})$  so  $Z_1, Z_2 \in \binom{X}{k}$  and  $|Z_2 \cap Y^*| = |Z_1 \cap Y^*|$  and  $Z_1 \cap \{a_{Y^*}, b_{Y^*}\} = Z_2 \cap \{a_{Y^*}, b_{Y^*}\}$  hence by 6.12(4) we have  $Z_2 \in \mathcal{P}_\pm$  and clearly  $|Z_1 \setminus Z_2| = 1$ .

By 6.15 we have  $\{a_{Z_1}, b_{Z_1}\} = \{a_{Z_2}, b_{Z_2}\}$ . Similarly by (\*) we can prove by induction on  $m = |Z \setminus Z_1|$  that  $\{a_{Z_1}, b_{Z_1}\} \subseteq Z \in \binom{X}{k} \Rightarrow Z \in \mathcal{P}_\pm$  &  $\{a_{Z_1}, b_{Z_1}\} = \{a_{Z_1}, b_{Z_1}\}$ . If ( $\beta$ ) of (c) fails, then for some  $Z_3 \in \mathcal{P}_\pm, \{a_{Z_1}, b_{Z_1}\} \not\subseteq Z_3$ . Easily  $\{a_{Z_3}, b_{Z_3}\} \subseteq Z \in \binom{X}{k} \Rightarrow Z \in \mathcal{P}_\pm$  &  $\{a_2, b_2\} = \{a_{Z_3}, b_{Z_3}\}$ . We can find  $Y \in \binom{X}{k}$ , such that  $\{a_{Z_1}, b_{Z_1}, a_{Z_3}, b_{Z_3}\} \subseteq Y$ ; contradiction.  $\square_{6.18}$

**6.18 Claim.** 1) The  $(*)$  of 5.6 holds.  
 2) In 6.17 clause (c), clause  $(\alpha)$  holds.

*Proof.* 1) Obvious by part (2) from  $(\alpha)$ .

2) First assume  $(\beta)$ , so by ?, clause (b), 6.3 + 6.4 we have without loss of generality  
 $\rightarrow$  scite{6.17} undefined

either  $\{a, b\} = \{a_{Y^*}, b_{Y^*}\} \subseteq Y \in \binom{X}{k} \Rightarrow f_Y(a, b) = b$  or  $\{a_{Y^*}, b_{Y^*}\} \subseteq Y \in \binom{X}{K} \Rightarrow f_Y(a_{Y^*}, b_{Y^*}) = b_{Y^*} = f(b_{Y^*}, a_{Y^*})$ . In both cases  $f^*$  is simple and not a monarchy contradiction to 5.2.

Second, assume  $(\gamma)$ . Let  $\langle \pi_i : i < i^* \rangle$  be a list of the permutations  $\pi$  of  $X$  such that  $\pi\{a_{Y^*}\} = \{a_{Y^*}, b_{Y^*}\}$ .

Let  $f_i^*$  be  $f^*$  conjugated by  $\pi_i$ . Now define  $g^i$  for  $i \leq i^*$  by induction on  $i$  :  
 $g_Y^0(x_1, x_2) = x_1, g_Y^{i+1}(x_1, x_2) = f_i^*(g_Y^i(x_1, x_2), x_2)$ . So  $g^{i^*} \in \mathcal{F}_{[2]}$  and  $\text{dom}_2(g^{i^*}) = \bigcap_{i < i^*} \text{dom}_2(f_i^*)$  where  $\text{dom}_2(g) = \{(Z, a, b) : a, b \in Z \in \binom{X}{k} \text{ and } g_Z(a, b) = b \neq a\}$ ,

so  $\text{dom}_1(g^{i^*}) = \bigcup_{i < i^*} \text{dom}_1(f_i^*)$  hence

$$(*)_1 \quad g_Y^{i^*}(a_1, a_2) = a_2 \text{ if } \{a_1, a_2\} = \{a_{Y^*}, b_{Y^*}\}$$

$$(*)_2 \quad g_Y^{i^*}(a_1, a_2) = a_1 \text{ if } \{a_1, a_2\} \neq \{a_{Y^*}, b_{Y^*}\}.$$

Now  $g$  is simple but not a monarchical contradiction to 5.2.

$\square_{6.18}$

§7 THE CASE  $r \geq 4$ 

7.1 Hypothesis.  $r = r(\mathcal{F}) \geq 4$ .

**7.2 Claim.** 1) For every  $f \in \mathcal{F}_r$  there is  $\ell(f) \in \{1, \dots, r\}$  such that

⊗ if  $Y \in \binom{X}{k}$ ,  $\bar{a} \in {}^r Y$ ,  $|Rang(\bar{a})| < r$  (i.e.  $\bar{a}$  is not one-to-one) then  $f_Y(\bar{a}) = a_{\ell(*)}$ .

2)  $r \leq k$ .

*Proof.* 1) Clearly there is a two-place function  $h$  from  $\{1, \dots, r\}$  to  $\{1, \dots, r\}$  such that: if  $y_1, \dots, y_r \in Y \in \binom{X}{k}$ ,  $y_\ell = y_k \wedge \ell \neq k \Rightarrow f_Y(y_1, \dots, y_r) = y_{h(\ell, k)}$ ; we have some freedom so let without loss of generality

$$\boxtimes \ell \neq k \Rightarrow h(\ell, k) \neq k.$$

Case 1: For some  $\bar{x} \in {}^r Y$ ,  $Y \in \binom{X}{k}$  and  $\ell_1 \neq k_1 \in \{1, \dots, r\}$  we have

$$|Rang(\bar{x})| = r - 1$$

$$x_{\ell_1} = x_{k_1}$$

$$f_Y(\bar{x}) \neq x_{\ell_1},$$

equivalently:  $h(\ell_1, k_1) \notin \{\ell, k\}$ .

Without loss of generality  $\ell_1 = r - 1, k_1 = r, f_Y(\bar{x}) = x_1$  (as by a permutation  $\sigma$  of  $\{1, \dots, r\}$  we can replace  $f$  by  $f^\sigma : f_Y^\sigma(x_1, \dots, x_r) = f_Y(x_{\sigma(1)}, \dots, x_{\sigma(r)})$ ).

We can choose  $Y \in \binom{X}{k}$  and  $x \neq y$  in  $Y$  so  $h(x, y, \dots, y) = x$  hence  $\ell \neq k \in \{2, \dots, r\} \Rightarrow h(\ell, k) = 1$ .

Now for  $\ell \in \{2, \dots, r\}$  we have agreed  $h(1, \ell) \neq \ell$ , (see  $\boxtimes$ ), as  $h \upharpoonright \{(m, n) : 1 \leq m < n \leq r\}$  is not constantly 1 (as  $f$  is not a monarchy) for some such  $\ell, h(1, \ell) \neq 1$  so without loss of generality  $h(1, 2) \neq 1, 2$ , so without loss of generality  $h(1, 2) = 3$  but as  $r \geq 4$  we have if  $x \neq y \in Y \in \binom{X}{k}$  then  $f_Y(x, x, y, y, \dots, y)$  is  $y$  as  $h(1, 2) = 3$  and is  $x$  as  $h(3, 4) = 1$ , contradiction. So

⊗  $h \upharpoonright \{(\ell, k) : 1 \leq \ell < k \leq r\}$  is constantly 1.

$$\bar{x} \in {}^r X \text{ is with repetitions} \Rightarrow h(\bar{x}) = x_1$$

as required.

Case 2: Not Case 1.

So  $h(\ell, k) \in \{\ell, k\}$  for  $\ell \neq k \in \{1, \dots, r\}$ . Now let  $Y \in \binom{X}{k}$ ,  $x \neq y \in Y$  and look at  $f_Y(x, x, y, y, \dots)$  it is both  $x$  as  $h(1, 2) \in \{1, 2\}$  and  $y$  as  $h(3, 4) \in \{3, 4\}$ , contradiction.

2) This follows as if  $f \in \mathcal{F}_{[r]}$  and  $k < r(\mathcal{F})$  and  $\ell(*)$  is as in part (1) then  $f_Y(\bar{x}) = x_{\ell(*)}$  always, as  $x_{\ell(*)}$  has repetitions by pigeon-hull.  $\square_{7.2}$

**7.3 Definition.**  $f = f_{r;\ell,k} = f^{r;\ell,k}$  is the  $r$ -place function

$$f_Y(\bar{x}) = \begin{cases} x_\ell & \bar{x} \text{ is with repetitions} \\ x_k & \text{otherwise} \end{cases}$$

**7.4 Claim.** 1) If  $f_{r;1,2} \in \mathcal{F}$  then  $f_{r;\ell,k} \in \mathcal{F}$  for  $\ell \neq k$ .

2) If  $f_{r;1,2} \in \mathcal{F}$ ,  $r \geq 3$  then  $f_{r+1;1,2} \in \mathcal{F}$ .

*Proof.* 1) Trivial.

2) Let  $r \geq 5$  for  $g(x_1, \dots, x_{r+1}) = f_{r;1,2}(x_1, x_2, \tau_3(x_1, \dots) \dots \tau_r(x_1, \dots))$  where  $\tau_m \equiv f_{r,1,m}(x_1, \dots, x_m, x_{m+2}, \dots, x_{r+1})$  that is  $x_{m+1}$  is omitted.

Continue as in the proof of 2.7.  $\square_{7.4}$

**7.5 Claim.** Assume  $Y \in \binom{X}{k}$ ,  $\bar{a} \in {}^r Y$  is one-to-one. There is  $f = f_{Y,\bar{a}} \in \mathcal{F}_r$  such that  $f_{Y,\bar{a}}(\bar{a}) = a_2$  and  $f_{Z,\bar{a}}(\bar{b}) = b_1$  if  $Z \in \binom{X}{k}$  and  $\bar{b} \in {}^r X$  is not one to one (so  $(*)$  of 5.6 holds).

*Proof.* Let  $f \in \mathcal{F}_r$  be not monarchical, and without loss of generality  $\ell(*) = 1$  in 7.2. By being not a monarchy, for some  $Y, \bar{a}$  and some  $k \in \{2, \dots, r\}$  we have  $f(\bar{a}) = a_k \neq a_1$ ; necessarily  $\bar{a}$  is one-to-one. Conjugating by  $\pi \in \text{Per}(X)$  we get any  $Y, \bar{a}$  and permuting  $[2, r]$ , we get  $f_{Y,\bar{a}}(\bar{a}) = a_2$ .  $\square_{7.3}$

## REFERENCES.

- [Arr50] Kenneth Arrow. A difficulty in the theory of social welfare. *J. of Political Economy*, **58**:328–346, 1950.
- [Fis73] Peter Fishburn. *The Theory of Social Choice*. Princeton University Press, 1973.
- [Kal01] Gil Kalai. Learnability and rationality of choice. 2001.
- [KRS01] Gil Kalai, Ariel Rubinstein, and Ran Spiegler. Comments on rationalizing choice functions which violate rationality. *Preprint*, 2001.
- [Pel84] Bezalel Peleg. *Game theoretic analysis of voting in committees*, volume 7 of *Econometric Society Monographs in Pure Theory*. Cambridge University Press, Cambridge, 1984.
- [RF86] Ariel Rubinstein and Peter Fishburn. Algebraic aggregation theory. *J. Econom. Theory*, **38**, 1986.
- [Sen86] Amartya Sen. Social choice theory. In Arrow and Intriligator, editors, *Handbook of Mathematical Economics, Vol III*, chapter 22, pages 1073–1182. North-Holland, Amsterdam, 1986.
- [Szb99] László Szabó. Algebras that are simple with weak automorphisms. *Algebra Universalis*, **42**:205–233, 1999.
- [Szn96] Ágnes Szendrei. Clones in universal algebra. volume 99 of *Séminaire de Mathématiques Supérieures [Seminar on Higher Mathematics]*. Presses de l’Université de Montréal, Montréal, QC, 1986.